

Marcus Fernando da Silva Praxedes
(Organizador)

Ciências da saúde:

Políticas públicas, assistência e gestão

Atena
Editora
Ano 2021

Marcus Fernando da Silva Praxedes
(Organizador)

Ciências da saúde:

Políticas públicas, assistência e gestão

Atena
Editora
Ano 2021

Editora chefe

Profª Drª Antonella Carvalho de Oliveira

Editora executiva

Natalia Oliveira

Assistente editorial

Flávia Roberta Barão

Bibliotecária

Janaina Ramos

Projeto gráfico

Camila Alves de Cremo

Daphynny Pamplona

Gabriel Motomu Teshima

Luiza Alves Batista

Natália Sandrini de Azevedo

Imagens da capa

iStock

Edição de arte

Luiza Alves Batista

2021 by Atena Editora

Copyright © Atena Editora

Copyright do texto © 2021 Os autores

Copyright da edição © 2021 Atena Editora

Direitos para esta edição cedidos à Atena Editora pelos autores.

Open access publication by Atena Editora



Todo o conteúdo deste livro está licenciado sob uma Licença de Atribuição Creative Commons. Atribuição-Não-Comercial-NãoDerivativos 4.0 Internacional (CC BY-NC-ND 4.0).

O conteúdo dos artigos e seus dados em sua forma, correção e confiabilidade são de responsabilidade exclusiva dos autores, inclusive não representam necessariamente a posição oficial da Atena Editora. Permitido o *download* da obra e o compartilhamento desde que sejam atribuídos créditos aos autores, mas sem a possibilidade de alterá-la de nenhuma forma ou utilizá-la para fins comerciais.

Todos os manuscritos foram previamente submetidos à avaliação cega pelos pares, membros do Conselho Editorial desta Editora, tendo sido aprovados para a publicação com base em critérios de neutralidade e imparcialidade acadêmica.

A Atena Editora é comprometida em garantir a integridade editorial em todas as etapas do processo de publicação, evitando plágio, dados ou resultados fraudulentos e impedindo que interesses financeiros comprometam os padrões éticos da publicação. Situações suspeitas de má conduta científica serão investigadas sob o mais alto padrão de rigor acadêmico e ético.

Conselho Editorial**Ciências Biológicas e da Saúde**

Prof. Dr. André Ribeiro da Silva – Universidade de Brasília

Profª Drª Anelise Levay Murari – Universidade Federal de Pelotas

Prof. Dr. Benedito Rodrigues da Silva Neto – Universidade Federal de Goiás

Profª Drª Daniela Reis Joaquim de Freitas – Universidade Federal do Piauí

Profª Drª Débora Luana Ribeiro Pessoa – Universidade Federal do Maranhão

Prof. Dr. Douglas Siqueira de Almeida Chaves – Universidade Federal Rural do Rio de Janeiro



Prof. Dr. Edson da Silva – Universidade Federal dos Vales do Jequitinhonha e Mucuri
Prof^o Dr^a Elizabeth Cordeiro Fernandes – Faculdade Integrada Medicina
Prof^o Dr^a Eleuza Rodrigues Machado – Faculdade Anhanguera de Brasília
Prof^o Dr^a Elane Schwinden Prudêncio – Universidade Federal de Santa Catarina
Prof^o Dr^a Eysler Gonçalves Maia Brasil – Universidade da Integração Internacional da Lusofonia Afro-Brasileira
Prof. Dr. Ferlando Lima Santos – Universidade Federal do Recôncavo da Bahia
Prof^o Dr^a Fernanda Miguel de Andrade – Universidade Federal de Pernambuco
Prof. Dr. Fernando Mendes – Instituto Politécnico de Coimbra – Escola Superior de Saúde de Coimbra
Prof^o Dr^a Gabriela Vieira do Amaral – Universidade de Vassouras
Prof. Dr. Gianfábio Pimentel Franco – Universidade Federal de Santa Maria
Prof. Dr. Helio Franklin Rodrigues de Almeida – Universidade Federal de Rondônia
Prof^o Dr^a Iara Lúcia Tescarollo – Universidade São Francisco
Prof. Dr. Igor Luiz Vieira de Lima Santos – Universidade Federal de Campina Grande
Prof. Dr. Jefferson Thiago Souza – Universidade Estadual do Ceará
Prof. Dr. Jesus Rodrigues Lemos – Universidade Federal do Piauí
Prof. Dr. Jônatas de França Barros – Universidade Federal do Rio Grande do Norte
Prof. Dr. José Max Barbosa de Oliveira Junior – Universidade Federal do Oeste do Pará
Prof. Dr. Luís Paulo Souza e Souza – Universidade Federal do Amazonas
Prof^o Dr^a Magnólia de Araújo Campos – Universidade Federal de Campina Grande
Prof. Dr. Marcus Fernando da Silva Praxedes – Universidade Federal do Recôncavo da Bahia
Prof^o Dr^a Maria Tatiane Gonçalves Sá – Universidade do Estado do Pará
Prof^o Dr^a Mylena Andréa Oliveira Torres – Universidade Ceuma
Prof^o Dr^a Natiéli Piovesan – Instituto Federacl do Rio Grande do Norte
Prof. Dr. Paulo Inada – Universidade Estadual de Maringá
Prof. Dr. Rafael Henrique Silva – Hospital Universitário da Universidade Federal da Grande Dourados
Prof^o Dr^a Regiane Luz Carvalho – Centro Universitário das Faculdades Associadas de Ensino
Prof^o Dr^a Renata Mendes de Freitas – Universidade Federal de Juiz de Fora
Prof^o Dr^a Vanessa da Fontoura Custódio Monteiro – Universidade do Vale do Sapucaí
Prof^o Dr^a Vanessa Lima Gonçalves – Universidade Estadual de Ponta Grossa
Prof^o Dr^a Vanessa Bordin Viera – Universidade Federal de Campina Grande
Prof^o Dr^a Welma Emidio da Silva – Universidade Federal Rural de Pernambuco



Ciências da saúde: políticas públicas, assistência e gestão

Diagramação: Daphynny Pamplona
Correção: Yaidy Paola Martinez
Indexação: Amanda Kelly da Costa Veiga
Revisão: Os autores
Organizador: Marcus Fernando da Silva Praxedes

Dados Internacionais de Catalogação na Publicação (CIP)

C569 Ciências da saúde: políticas públicas, assistência e gestão /
Organizador Marcus Fernando da Silva Praxedes. –
Ponta Grossa - PR: Atena, 2021.

Formato: PDF

Requisitos de sistema: Adobe Acrobat Reader

Modo de acesso: World Wide Web

Inclui bibliografia

ISBN 978-65-5983-765-6

DOI: <https://doi.org/10.22533/at.ed.656211012>

1. Ciências da saúde. I. Praxedes, Marcus Fernando da
Silva (Organizador). II. Título.

CDD 613

Elaborado por Bibliotecária Janaina Ramos – CRB-8/9166

Atena Editora
Ponta Grossa – Paraná – Brasil
Telefone: +55 (42) 3323-5493
www.atenaeditora.com.br
contato@atenaeditora.com.br



DECLARAÇÃO DOS AUTORES

Os autores desta obra: 1. Atestam não possuir qualquer interesse comercial que constitua um conflito de interesses em relação ao artigo científico publicado; 2. Declaram que participaram ativamente da construção dos respectivos manuscritos, preferencialmente na: a) Concepção do estudo, e/ou aquisição de dados, e/ou análise e interpretação de dados; b) Elaboração do artigo ou revisão com vistas a tornar o material intelectualmente relevante; c) Aprovação final do manuscrito para submissão.; 3. Certificam que os artigos científicos publicados estão completamente isentos de dados e/ou resultados fraudulentos; 4. Confirmam a citação e a referência correta de todos os dados e de interpretações de dados de outras pesquisas; 5. Reconhecem terem informado todas as fontes de financiamento recebidas para a consecução da pesquisa; 6. Autorizam a edição da obra, que incluem os registros de ficha catalográfica, ISBN, DOI e demais indexadores, projeto visual e criação de capa, diagramação de miolo, assim como lançamento e divulgação da mesma conforme critérios da Atena Editora.



DECLARAÇÃO DA EDITORA

A Atena Editora declara, para os devidos fins de direito, que: 1. A presente publicação constitui apenas transferência temporária dos direitos autorais, direito sobre a publicação, inclusive não constitui responsabilidade solidária na criação dos manuscritos publicados, nos termos previstos na Lei sobre direitos autorais (Lei 9610/98), no art. 184 do Código Penal e no art. 927 do Código Civil; 2. Autoriza e incentiva os autores a assinarem contratos com repositórios institucionais, com fins exclusivos de divulgação da obra, desde que com o devido reconhecimento de autoria e edição e sem qualquer finalidade comercial; 3. Todos os e-book são *open access*, *desta forma* não os comercializa em seu site, sites parceiros, plataformas de *e-commerce*, ou qualquer outro meio virtual ou físico, portanto, está isenta de repasses de direitos autorais aos autores; 4. Todos os membros do conselho editorial são doutores e vinculados a instituições de ensino superior públicas, conforme recomendação da CAPES para obtenção do Qualis livro; 5. Não cede, comercializa ou autoriza a utilização dos nomes e e-mails dos autores, bem como nenhum outro dado dos mesmos, para qualquer finalidade que não o escopo da divulgação desta obra.



APRESENTAÇÃO

Apresentamos a coleção “Ciências da Saúde: Políticas Públicas, Assistência e Gestão”, que reúne trabalhos científicos relevantes das mais diversas áreas da Ciência da Saúde. A coleção divide-se em dois volumes, em que o objetivo central foi apresentar de forma categorizada e clara estudos desenvolvidos em diversas instituições de ensino e pesquisa nacionais e internacionais.

O primeiro volume traz estudos relacionados à assistência em saúde com abordagem do uso correto dos medicamentos, com uma discussão relevante sobre a automedicação e adesão ao tratamento, bem como da importância de uma abordagem interprofissional; uso de fitoterápicos; alimentação saudável; segurança do paciente e qualidade do cuidado; assistência em saúde no domicílio e uso de ferramentas para avaliação em saúde.

O segundo volume reúne variados estudos que abordam temáticas que continuam atuais e sensíveis às políticas públicas e para uma melhor gestão em saúde. Dentre algumas discussões, tem-se a violência contra a mulher e a necessidade do empoderamento feminino, bem como da adequada assistência às vítimas; questões psicossociais; o uso de tecnologias em saúde; abordagem de doenças negligenciadas; qualidade da água e de alimentos consumidos pela população; a importância da auditoria em saúde, do planejamento estratégico e da importância da capacitação profissional para o exercício da gestão em saúde.

Espera-se que os trabalhos científicos apresentados possam servir de base para uma melhor assistência, gestão em saúde e desenvolvimento de políticas públicas mais efetivas. Nesse sentido, a Atena Editora se destaca por possuir a estrutura capaz de oferecer uma plataforma consolidada e confiável para estes pesquisadores exporem e divulguem seus resultados.


Marcus Fernando da Silva Praxedes

SUMÁRIO

CAPÍTULO 1..... 1

AUTOMEDICAÇÃO EM PACIENTES IDOSOS


Katia Carvalho Marques
Ladislau Henrique Macedo dos Santos
Lucilene Carvalho Marques
Anna Maly Leão Neves Eduardo

 <https://doi.org/10.22533/at.ed.6562110121>

CAPÍTULO 2..... 12

AUTOMEDICAÇÃO ENTRE PROFISSIONAIS DA ENFERMAGEM E OS FATORES QUE FAVORECEM ESSA PRÁTICA EM UM HOSPITAL PÚBLICO DE MANAUS - AM


Adriane Kakijima Bonfim
Geliane da Gama Lima Torres
Liliane Íris Bonfim Pinheiro
Mychele Azevedo Lima
Silas Pereira Muraiare
Leslie Bezerra Monteiro
Silvana Nunes Figueiredo
Hanna Lorena Morais Gomes
Andreia Silvana Silva Costa
Loren Anselmo do Nascimento

 <https://doi.org/10.22533/at.ed.6562110122>

CAPÍTULO 3..... 24

PARTICIPAÇÃO ATIVA DO FARMACÊUTICO FRENTE À AUTOMEDICAÇÃO


Luiz Olivier Rocha Vieira Gomes
Nadyellem Graciano da Silva
Simone Soares da Silva
Axell Donelli Leopoldino Lima
Ivone Oliveira da Silva
Mônica Larissa Gonçalves da Silva
Elizabeth Moreira Klein
Rodrigo Lima dos Santos Pereira
Victória Melo da Costa
Paulo Diniz de Oliveira
Andréa Fernanda Luna Rodrigues
Eduarda Rocha Teixeira Magalhães
Lustarlone Bento de Oliveira

 <https://doi.org/10.22533/at.ed.6562110123>

CAPÍTULO 4..... 38

ATENÇÃO FARMACÊUTICA NO USO INADEQUADO DE PSICOTRÓPICOS


Lucimara Regina Aleixo Ferreira
Maria Adellane de Oliveira Silva
Heleneide Cristina Campos Brum

 <https://doi.org/10.22533/at.ed.6562110124>

CAPÍTULO 5..... 51

ESTIMATIVA DE ADESÃO A MEDICAMENTO ANTIRRETROVIRAL COFORMULADO


Yanna Dantas Rattmann
Bárbara Thaís Polisel de Sá
Mariana Ribeiro Martins
Leticia Mara Marca
Débora Bauer Schultz
Flavia Helen Correia
Sacha Testoni Lange
Marina Yoshie Miyamoto
Beatriz Böger
Frederico Alves Dias

 <https://doi.org/10.22533/at.ed.6562110125>

CAPÍTULO 6..... 61

INCONFORMIDADES RELACIONADAS À UTILIZAÇÃO DE MEDICAMENTOS ORAIS ADMINISTRADOS VIA SONDAS EM UMA INSTITUIÇÃO HOSPITALAR


Sílvia Maria Jacques Neves
Andreia Insabralde de Queiroz Cardoso
Ramon Moraes Penha
Elza Aparecida Machado Domingues
Camila Guimarães Polisel

 <https://doi.org/10.22533/at.ed.6562110126>

CAPÍTULO 7..... 77

A ATUAÇÃO DO ENFERMEIRO NA ÁREA DE QUIMIOTERAPIA E OS RISCOS OCUPACIONAIS

Fernanda da Silva Ferreira
Larissa Bartles dos Santos
Stefany Pinheiro de Moura
Rutiana Santos Batista
Gilvania Santos Ferreira Sousa
Tatiane Regina de Souza Castro
Mariana Machado Figueiredo
Bernadete de Lourdes Xavier
Maria Gabriela Lourenço
Tássara Vitória da Silva Almeida
Maria Eduarda Pinto Pinheiro
Letícia F. Fiuza Bacelar

 <https://doi.org/10.22533/at.ed.6562110127>


CAPÍTULO 8..... 86

CONTRACEPTIVO DE EMERGÊNCIA

Alex Sandro Pereira Ivasse
Benjamim De Almeida Silva

Paulo Roberto De Sousa Lima Junior

Anna Maly Leão Neves Eduardo

 <https://doi.org/10.22533/at.ed.6562110128>

CAPÍTULO 9..... 95


FITOTERÁPICOS E PLANTAS MEDICINAIS NO BRASIL: UMA ABORDAGEM SOBRE LEGISLAÇÃO NO CONTEXTO DAS POLÍTICAS DE SAÚDE

Muiara Aparecida Moraes

Aílson da Luz André de Araújo

Ana Lúcia Santos de Matos Araújo

Orlando Vieira de Sousa

 <https://doi.org/10.22533/at.ed.6562110129>

CAPÍTULO 10..... 109

PALMÁCEAS REGIONAIS: UMA REVISÃO EDUCATIVA DA IMPORTÂNCIA NA PRODUÇÃO DE LIPÍDIOS E APLICAÇÃO SUSTENTÁVEL EM PRODUTOS PARA SAÚDE

Rafael Miranda Carvalho Dos Reis


Vitória Ellen Batista de Moraes Nascimento

Alana Oliveira de Sena

Leidiane Rodrigues Santiago Feitosa

Leonardo Fonseca Maciel

Neila de Paula Pereira

 <https://doi.org/10.22533/at.ed.65621101210>

CAPÍTULO 11..... 130


A EFICÁCIA DO CANABIDIOL NO TRATAMENTO DE EPILEPSIA

Denisia verônica Pereira dos Santos

Larissa Aparecida Alves Ferreira

Lucas Cardoso Lopes

Anna Maly de Leão e Neves Eduardo

 <https://doi.org/10.22533/at.ed.65621101211>

CAPÍTULO 12..... 137

FACTORES ASOCIADOS A LA PÉRDIDA DE PESO DE LOS PACIENTES Y LA DIETA PRESCRITA DURANTE LA HOSPITALIZACIÓN

Vânia Aparecida Leandro-Merhi


José Luis Braga de Aquino

Hallan Douglas Bertelli

Geovanna Godoy Ramos

Elisa Teixeira Mendes

José Alexandre Mendonça


 <https://doi.org/10.22533/at.ed.65621101212>

CAPÍTULO 13..... 153

CAPACIDADE DO CHÁ VERDE NO AUXÍLIO DO TRATAMENTO DA OBESIDADE BEM COMO DE SUAS COMORBIDADES (UMA REVISÃO)

Débora Gracielly da Silva

Maria José Arruda De Albuquerque Lopes
Raquel Maria da Silva
Jobson Josimar Marques Teixeira
José Edson de Souza Silva

 <https://doi.org/10.22533/at.ed.65621101213>

CAPÍTULO 14..... 162

ASSOCIAÇÃO ENTRE PERDA AUDITIVA E ALIMENTAÇÃO: PROMOÇÃO DA SAÚDE


Patrícia Haas
Laura Faustino Gonçalves
Beatriz Vitorio Ymai Rosendo
Karina Mary Paiva
Rodrigo Sudatti Delevatti

 <https://doi.org/10.22533/at.ed.65621101214>

CAPÍTULO 15..... 178

**A SEGURANÇA DO PACIENTE INSERIDA NA GESTÃO DA QUALIDADE HOSPITALAR:
UMA PROPOSTA SIMPLIFICADA DE IMPLANTAÇÃO**


Fabiano Lucio de Almeida Silva

 <https://doi.org/10.22533/at.ed.65621101215>

CAPÍTULO 16..... 191

**INTERFERÊNCIA DO DIABETES *Mellitus* NA SAÚDE NUTRICIONAL DE PESSOAS
IDOSAS**

Carina Barbosa Bandeira
Maria Vieira de Lima Saintrain
Rafaela Laís e Silva Pesenti Sandrin
Marina Arrais Nobre
Ana Ofélia Lima Portela
Debora Rosana Alves Braga de Figueiredo
Maria da Glória Almeida Martins
Maria Isabel Damasceno Martins Fernandes
Camila Bandeira de Sousa
Anna Cecília Nunes dos Santos
Janaína Alvarenga Aragão
Luciano Silva Figueiredo


 <https://doi.org/10.22533/at.ed.65621101216>

CAPÍTULO 17..... 202

**PANORAMA GERAL SOBRE AS COMPETÊNCIAS E PRÁTICAS PROFISSIONAIS NA
QUALIDADE E SEGURANÇA NO CUIDADO DO PACIENTE**

Francisca das Chagas Sheyla Almeida Gomes Braga
Cléciton Braga Tavares
Geisa Machado Fontenelle
Roxana Mesquita de Oliveira Teixeira Siqueira
Antônio Francisco Machado Pereira
Yara Maria Rêgo Leite


Veronica Elis de Araújo Rezende
Adriana Jorge Brandão
Maria Lailda de Assis Santos
Sandra Valéria Nunes Barbosa
Luciane Resende da Silva Leonel

 <https://doi.org/10.22533/at.ed.65621101217>

CAPÍTULO 18.....210

O CUIDADO DOMICILIAR COMO ESTRATÉGIA DE ASSISTÊNCIA NO PERÍODO DA PANDEMIA (COVID 19)

Camila Augusta de Oliveira Sá
Diana Muniz Pinto
Lúcia Helena Gonçalves Martins
Mariana Freitas e Silva Maia
Ney Sindeaux Moreira

 <https://doi.org/10.22533/at.ed.65621101218>

CAPÍTULO 19.....217

SOBRECARGA DE CUIDADOS DECORRENTE DA DOENÇA DE ALZHEIMER: UMA REVISÃO DE LITERATURA


Mari Ângela Victoria Lourenci Alves
Aline dos Santos Duarte
Bibiana Fernandes Trevisan
Michelle Batista Ferreira
Rodrigo D Ávila Lauer
Tábata de Cavata Souza

 <https://doi.org/10.22533/at.ed.65621101219>

CAPÍTULO 20.....223

VISITA DOMICILIAR COMO MECANISMO DE ATUAÇÃO DE RESIDENTES EM ATENÇÃO HOSPITALAR À SAÚDE: RELATO DE EXPERIÊNCIA

Danielle Ramos Domenis
Janayna de Almeida Andrade
Ranna Adrielle Lima Santos
Suzanne Guimarães Machado
Felipe Douglas Silva Barbosa


 <https://doi.org/10.22533/at.ed.65621101220>

CAPÍTULO 21.....232

PRIMEIROS SOCORROS PARA LEIGOS: EXPERIÊNCIA DE EXTENSÃO UNIVERSITÁRIA MEDIADA PELA INTERNET

Paulo Cilas de Carvalho Sousa
Jaqueline Renata da Silva Brito
Fernanda Karielle Coelho Macedo
Maria Eduarda de Sousa Brito
Oyama Siqueira Oliveira
Lairton Batista de Oliveira

Francisco Gilberto Fernandes Pereira


 <https://doi.org/10.22533/at.ed.65621101221>

CAPÍTULO 22.....241

PERFIL DE PACIENTES ATENDIDOS NO AMBULATÓRIO DE ENDOCRINOLOGIA PEDIÁTRICA EM UM HOSPITAL UNIVERSITÁRIO

Andréa Timóteo dos Santos Dec

Margarete Aparecida Salina Maciel

 <https://doi.org/10.22533/at.ed.65621101222>

CAPÍTULO 23.....255

CARACTERIZAÇÃO DO PERFIL, DA PERCEPÇÃO DO AMBIENTE OCUPACIONAL E DOS PRINCIPAIS DESFECHOS OSTEOMUSCULARES NOS FUNCIONÁRIOS DA ATENÇÃO BÁSICA A SAÚDE DO MUNICÍPIO DE PRESIDENTE KUBITSCHECK – MINAS GERAIS

Alysson Geraldo Mendonça

 <https://doi.org/10.22533/at.ed.65621101223>

CAPÍTULO 24.....266

APLICAÇÃO DA ESCALA BIANCHI DE STRESS EM BLOCO OPERATÓRIO

Maria Helane Rocha Batista Gonçalves

Christian Raphael Fernandes Almeida

Kelly Barros Marques

Rafaella Regis de Albuquerque Isacksson

Débora Rodrigues Guerra Probo

 <https://doi.org/10.22533/at.ed.65621101224>

CAPÍTULO 25.....279

USO DE QUESTIONÁRIOS COMO FERRAMENTAS PARA AVALIAÇÃO DE DISBIOSE INTESTINAL E RISCO PARA DOENÇAS CRÔNICAS NÃO TRANSMISSÍVEIS: UMA REVISÃO DE LITERATURA

Whellyda Katrynne Silva Oliveira

Débora Paloma de Paiva Sousa

Heide Sara Santos Ferreira

Vitória Ribeiro Mendes

Lana Maria Mendes Gaspar

Joyce Sousa Aquino Brito

Andressa Correia das Neves

Juliana Feitosa Ferreira

Elinayara Pereira da Silva

Marta Gama Marques Castro

Vanessa Gomes de Oliveira

Stefany Rodrigues de Sousa Melo

 <https://doi.org/10.22533/at.ed.65621101225>

CAPÍTULO 26.....289


TRATAMENTO FARMACOLÓGICO DO DESEJO SEXUAL HIPOATIVO FEMININO:

MEDICAÇÕES APROVADAS PELO FOOD AND DRUG ADMINISTRATION (FDA)

Gabriela Pascueto Amaral

Nathalie de Paula Damião

Lúcio Mauro Bisinotto Júnior

 <https://doi.org/10.22533/at.ed.65621101226>

CAPÍTULO 27.....299

OS PRINCIPAIS IMPACTOS À SAÚDE DA CRIANÇA CAUSADOS PELO CONSUMO DE CORANTES ALIMENTÍCIOS: UMA REVISÃO BIBLIOGRÁFICA

Marcelo Borges Figueira da Mota

Brunna Michelly da Silva Sousa

Tamyres Borges Pereira

Isabella Chaves Lira Cruz

Juliana Amorim Alfaix Natário

Irlane Moraes Vasconcelos Souza


Antonina Linhares Moraes Neta

Guilherme de Souza Gomes

Fernanda de Melo Franco Machado

Enzo Cardoso de Faria

Gabriel Mazuchini Belai

 <https://doi.org/10.22533/at.ed.65621101227>

SOBRE O ORGANIZADOR.....308

ÍNDICE REMISSIVO.....309

FACTORES ASOCIADOS A LA PÉRDIDA DE PESO DE LOS PACIENTES Y LA DIETA PRESCRITA DURANTE LA HOSPITALIZACIÓN

Data de aceite: 01/11/2021

Vânia Aparecida Leandro-Merhi

Postgraduate Program in Health Sciences.
Pontifícia Universidade Católica de Campinas
Campinas, São Paulo. Brazil

José Luis Braga de Aquino

Postgraduate Program in Health Sciences.
Pontifícia Universidade Católica de Campinas
Campinas, São Paulo. Brazil

Hallan Douglas Bertelli

Postgraduate Program in Health Sciences.
Pontifícia Universidade Católica de Campinas
Campinas, São Paulo. Brazil

Geovanna Godoy Ramos

Scholarship of Scientific Initiation. Faculdade
de Nutrição. Pontifícia Universidade Católica de
Campinas
Campinas, São Paulo. Brazil

Elisa Teixeira Mendes

Postgraduate Program in Health Sciences.
Pontifícia Universidade Católica de Campinas
Campinas, São Paulo. Brazil

José Alexandre Mendonça

Postgraduate Program in Health Sciences
Pontifícia Universidade Católica de Campinas
Campinas, São Paulo. Brazil

RESUMO: OBJETIVO: Avaliar a perda de peso ,dieta prescrita, e o status nutricional em pacientes hospitalizados, bem como seus fatores associados; **MÉTODOS:** perda de peso

durante a hospitalização, status nutricional, tipo de doença, e dieta prescrita foram investigados em um estudo retrospectivo em 621 pacientes hospitalizados. Os testes de Qui-quadrado, Fisher, Mann- Whitney e Kruskal-Wallis foram usados para análise estatística. Para identificação dos fatores associados com perda de peso, foi utilizado análise de regressão logística. O nível de significância adotado foi de 5%; **RESULTADOS:** Os pacientes com perda de peso durante a hospitalização foram associados com maior tempo de internação ($p < 0,0001$; OR=1.052; 95% CI= 1.030 a 1.073), desnutrição pela avaliação global subjetiva ($p = 0,0081$; OR=3.177; 95% CI=1.351 a 7.469) e neoplasias digestivas ($p = 0,0407$; OR=2.410; 95% CI=1.038 a 5.597); **CONCLUSÃO:** Perda de peso durante a hospitalização foi associado com neoplasias, doenças digestivas, desnutrição e maior tempo de internação.

PALAVRAS-CHAVE: Perda de peso. Dieta prescrita. Pacientes hospitalizados. Fatores associados

FACTORS ASSOCIATED WITH PATIENT WEIGHT LOSS AND PRESCRIBED DIET DURING HOSPITALIZATION

ABSTRACT: Objective: the aim of this study was to assess weight loss, diet prescribed, and nutritional status in hospitalized patients, as well as their associated factors. **Methods:** weight loss during hospitalization, nutritional status, disease type, and prescribed diet were investigated in a retrospective study in 621 hospitalized patients. The chi-squared, Fisher's, Mann-Whitney, and Kruskal-Wallis tests were used for statistical

analysis. To identify factors associated with weight loss a logistic regression analysis was performed. The significance level adopted for statistical tests was 5 %. **Results:** patients who experienced weight loss during hospitalization were associated with longer hospital stays ($p < 0.0001$; OR = 1.052; 95 % CI = 1.030 to 1.073), malnourishment according to the subjective global assessment ($p = 0.0358$; OR = 1.520; 95 % CI = 1,028 to 2,248), digestive disorders ($p = 0.0081$; OR = 3.177; 95 % CI = 1.351 to 7.469), and digestive neoplasms ($p = 0.0407$; OR = 2.410; 95 % CI = 1.038 to 5.597). **Conclusion:** weight loss during hospitalization was associated with neoplasms, digestive diseases, malnutrition, and length of stay.

KEYWORDS: Weight loss. Diet prescribed. Hospitalized patients. Factors associated.

This work was presented as a poster at the ESPEN 2020 Virtual Congress event.

Statement of authorship: VALM and JLBA equally contributed to the conception and design of the research. JAM contributed to the design of the research. VALM, HDB, GGR, and JLBA contributed to the acquisition and analysis of the data. ETM and JAM contributed to the interpretation of the data. VALM, JLBA, ETM, and JAM drafted the manuscript. All authors critically revised the manuscript, agree to be fully accountable for ensuring the integrity and accuracy of the work, and read and approved the final manuscript.

RESUMEN: Objetivo: el objetivo de este estudio fue evaluar la pérdida de peso, la dieta prescrita y el estado nutricional de pacientes hospitalizados y sus factores asociados.

Métodos: se investigó la pérdida de peso durante la hospitalización, el estado nutricional, el tipo de enfermedad y la dieta prescrita en un estudio retrospectivo de 621 pacientes hospitalizados. Las pruebas del chi cuadrado, Fisher, Mann-Whitney y Kruskal-Wallis se utilizaron para el análisis estadístico. Para identificar los factores asociados con la pérdida de peso se utilizó la regresión logística. El nivel de significación adoptado para las pruebas estadísticas fue del 5 %. **Resultados:** los casos de pérdida de peso durante la hospitalización se asociaron a las estancias hospitalarias más largas ($p < 0,0001$; OR = 1,052; IC 95 % = 1,030; 1,073), la desnutrición según la evaluación global subjetiva ($p = 0,0358$; OR = 1,520; IC 95 % = 1,028; 2,248) los trastornos digestivos ($p = 0,0081$; OR = 3,177; IC 95 % = 1,351; 7,469) y las neoplasias digestivas ($p = 0,0407$; OR = 2,410; IC 95 % = 1,038; 5,597).

Conclusión: la pérdida de peso durante la hospitalización se asoció con las neoplasias y las enfermedades digestivas, la desnutrición y la duración de la estancia.

PALABRAS CLAVE: Pérdida de peso. Dieta prescrita. Pacientes hospitalizados. Factores asociados.

INTRODUCTION

Several studies have investigated malnutrition (1-3) in the hospital setting, showing the impact of nutritional risk (2) on clinical outcome and hospital costs. Some investigations (3) have evaluated factors associated with malnutrition, pointing out the need for greater attention to nutritional care. Correia et al, in 2017 (1), showed that most studies describe a high prevalence of malnutrition, with rates ranging from 40 % to 60 %. The authors reported an increase in malnutrition associated with increased length of hospital stay, and with clinical and infectious complications. Thus, several nutritional indicators (4,5) have been used to assess nutritional status. Some (6) show that in individuals identified by risk screening as

being at nutritional risk, the diagnosis of malnutrition should be based on low body mass index or on combined weight loss indexes together with other indicators (6).

Nutritional intervention has contributed to reduce the risk of death and rehospitalization due to complications arising from poor nutritional status (7). In a prospective study (8), the priority of identifying nutritional status by means of nutritional screening instruments in clinical practice, in order to reduce morbidity and mortality rates (8), has been demonstrated. Other investigations have evidenced the importance of nutritional intervention with the use of specific supplements adapted to different clinical conditions (9). Another study (10) investigating nutritional status showed an association with age, nutritional therapy, and food intake, which were related to clinical outcomes such as prolonged hospital stay. Within this context, hospital weight loss, food consumption, and type of nutritional therapy (10-12) have been investigated in the relevant literature, and have motivated the performance of this study. Thus, the aim of this study was to assess weight loss, the diet prescribed to, and the nutritional status of hospitalized patients and their associated factors.

METHODS

STUDY DESIGN, ETHICAL APPROVAL AND POPULATION CHARACTERISTICS

This was a retrospective study with data collected from medical records at a university hospital, conducted between the years 2018 and 2019. The study was approved by the institution's Research and Ethics Committee. The inclusion criteria adopted were age equal to or greater than 20 years, investigation of nutritional status within 48 hours of hospitalization, and complete nutritional information in the hospital medical records. Patients with incomplete records, terminal illness, and hospitalized only for clinical investigation and/or exams were excluded.

It is worth mentioning that in this hospital department a routine nutritional status assessment is carried out at the very beginning of hospitalization, at most within two days of admission. This procedure is routinely performed in hospital clinical practice by a team of nutritionists, residents and interns trained in nutrition at this institution, who assessed the nutritional indicators of anthropometry, used laboratory tests, administered nutritional screening instruments, and reviewed food histories. These data are routinely recorded in the medical records of inpatients.

For the composition of the sample to be studied in this investigation, all patients admitted to surgery services who met all inclusion criteria in the period defined for the study were selected. Initially 673 patients were recruited for this study, but 52 of them did not meet all the inclusion criteria. Thus, 621 adult inpatients of both genders were considered eligible for the study.

METHODOLOGICAL PROCEDURES

Data concerning type of disease, length of stay, age, gender, recent weight loss, weight loss during hospitalization, subjective global assessment, nutritional risk screening, anthropometry, type of diet prescribed on admission, alteration and acceptance of diet, use of protein-calorie supplementation during hospitalization, and percentage of habitual energy intake in relation to the estimated energy requirements (%HEI/EER) are described below.

Recent weight loss (RWL)

RWL was considered unintentional when the patient reported having lost > 5 % of their weight within the last 3 months, prior to hospitalization (6).

Weight loss during hospitalization

In order to assess weight loss during the hospitalization period, the present study considered the loss of at least 1 kg of weight from admission (1st day of hospitalization) to hospital discharge (last day of hospitalization). This criterion was established by the study authors to take into account in-hospital weight loss.

Subjective global assessment (SGA)

Nutritional status was assessed by the SGA and was classified according to standard cutoff points (13) into well nourished and malnourished (in this study, patients classified as mildly, moderately, and severely malnourished were all considered malnourished).

Nutritional risk screening (NRS)

Nutritional risk was assessed using the NRS tool, classifying patients as with a score ≥ 3 (at risk) or < 3 (without risk) (14,15).

Anthropometry

Body mass index (BMI) was evaluated according to standard cutoff points for adults (16) and the elderly (17), and body composition parameters such as arm circumference (AC), triceps skinfold (TSF), and arm muscle circumference (AMC) (18,19).

Type of diet prescribed at admission, diet change, diet acceptance, and use of protein-calorie supplementation during hospitalization

The type of diet prescribed during hospitalization was evaluated and classified as general diet (with solid food), soft diet (with soft food), liquid diet (with liquid food), watery diet (water, tea and gelatin; this diet was used during short post-operative periods), enteral and parenteral diet, or fasting (oral fasting). We investigated whether there was any change

or alteration (yes/no) in the diet prescribed during hospitalization. For the evaluation of diet acceptance, acceptance was considered to be good when the patient reported consuming the totality of the diet offered; to be regular when the patient reported partial acceptance of the diet (not eating the entire diet offered); and to be poor when the patient had a low dietary intake. The use or not of calorie-protein nutritional supplements in liquid form (yes/no) during hospitalization was also surveyed.

Rate of habitual energy intake in relation to estimated energy requirements (%HEI/EER)

By nutritional anamnesis, the habitual energy intake of the week prior to hospitalization was evaluated through an individualized assessment of all patients, investigating their usual diet (recall of habitual consumption) and type and quantity of food, fractioning, and meal times. This allowed to calculate total energy consumption in relation to estimated energy requirements using the Harris & Benedict equation (20). Subsequently, the rate of energy intake in relation to energy requirements was considered in this study as being: $< 75\%$ or $\geq 75\%$ (15,21,22).

STATISTICAL ANALYSIS

Initially, a descriptive analysis of the data was performed with the presentation of frequency tables for categorical variables and measures of position and dispersion for continuous variables. Then, for comparison of proportions, a Chi-square test or Fisher's exact test was used when necessary. To compare continuous or orderable measurements between 2 groups, the Mann-Whitney test was applied; and to do so between 3 or more groups, the Kruskal-Wallis test, followed by Dunn's test to find the differences, when necessary, was used. Subsequently, a logistic regression analysis was used to identify factors associated with weight loss during hospitalization. The power of the study was calculated for the results of logistic regressions. Power ranged from 0.70 to > 0.90 . For the subjective global assessment (SGA) tool, power was 0.70 (70% power), and for the other variables it was > 0.90 ($> 90\%$). The level of significance adopted for the statistical tests was 5% (23-25).

RESULTS

The average age of the population studied was 57.85 ± 15.23 years, and the length of hospital stay was 10.80 ± 8.91 days. The general characteristics of the population are described in table I. When comparing the groups of patients with and without weight loss during hospitalization, there was a significant difference for length of stay ($p < 0.0001$), fasting time ($p = 0.0002$), SGA ($p = 0.0351$), and type of disease ($p = 0.0171$). In the study of factors associated with weight loss by logistic regression analysis, it was found that patients who experienced weight loss during hospitalization were associated with a longer hospital stay

($p < 0.0001$; OR = 1.052; 95 % CI = 1.030; 1.073), malnourishment by SGA ($p = 0.0358$; OR = 1.520; 95 % CI = 1.028; 2.248), digestive tract diseases (DTD) ($p = 0.0081$; OR = 3.177; 95 % CI = 1.351; 7.469), and digestive neoplasms ($p = 0.0407$; OR = 2.410; 95 % CI = 1.038; 5.597) (Table IIA). The other variables studied did not show a statistically significant difference (Tables IIA and IIB).

Table III shows a comparison between the types of diet prescribed during hospitalization and study variables. There was a statistically significant difference in the relationship between all study variables and type of diet prescribed during hospitalization. There was a significant difference between types of diet according to malnutrition classification by SGA, with a higher percentage of enteral and parenteral diet ($p < 0.0001$; chi-square test). There

Variables	Category	Frequency	Percentage
Gender	Female	227	36.55
	Male	394	63.45
Type of disease	Digestive tract disease	78	12.56
	Fractures and trauma	55	8.86
	Renal and urological	46	7.41
	Vascular	105	16.91
	Infections	19	3.06
	Head and neck neoplasms	48	7.73
	Neoplasms of the digestive tract	92	14.81
	Other diseases	100	16.10
	Other neoplasms	78	12.56
Hospitalization time	Up to 7 days	280	45.09
	> 7 days	341	54.91
Prescribed diet	Soft	144	23.30
	Enteral/Parenteral	37	5.99
	General	248	40.13
	Watery/Fasting	148	23.95
	Liquid	41	6.63
Supplement use	No	583	93.88
	Yes	38	6.12
Diet change	No	316	51.13
	Yes	302	48.87
Recent weight loss*	No	446	75.98
	Yes	141	24.02
Weight loss during hospitalization (≤ 1 kg)	No	436	70.21
	Yes	185	29.79

Global subjective assessment	Well nourished	474	76.33
	Malnourished	147	23.67
Nutritional risk screening	At risk	246	39.94
	No risk	370	60.06
Body mass index	Overweight	271	43.78
	Normal weight	250	40.39
	Low weight	98	15.83
%HEI/EER [†]	< 75 %	409	67.72
	≥ 75 %	195	32.28

Table I. General characteristics of the studied population (n = 621 patients)

*Recent weight loss (prior to hospitalization). [†]Percentage of habitual energy consumption in relation to estimated energy requirements.

was a significant difference between types of diet according to type of disease ($p < 0.0001$; Fisher's test). It was found that patients with digestive tract disease were prescribed fasting, water, and liquid diet with greater frequency. In cases of head and neck neoplasms and digestive tract neoplasms, a higher percentage of enteral and parenteral diets was observed (Table III).

Figure 1 shows the percentage distribution of the type of diet prescribed during hospitalization, according to weight loss (loss, or no loss, of at least 1 kg of weight during hospitalization). There was no statistically significant difference between patients who presented or did not present weight loss during hospitalization and the types of diets prescribed ($p = 0.1299$; chi-square test).

Variables	Category	No weight loss n = 436	With weight loss n = 185	p-value	p-value [†]	OR	95 % CI
Age	X ± SD	58.3 ± 15.3	56.8 ± 15.1	0.2469*	0.2654	0.994	0.983; 1.005
HT	X ± SD	9.6 ± 8.4	13.7 ± 9.3	<0.0001*	<0.0001	1.052	1.030; 1.073
FT	X ± SD	1.8 ± 2.5	2.2 ± 1.6	0.0002*	0.1599	1.109	0.960; 1.281
Gender, n (%)	Female	167 (38.3)	60 (32.4)	0.1648 [†]	0.1653	1.293	0.899; 1.860
	Male	269 (61.7)	125 (67.6)				
HT in days, n (%)	Up to 7 days	231 (53.0)	49 (26.5)	<0.0001 [†]	<0.0001	ref	ref
	> 7 days	205 (47.0)	136 (73.5)			3.128	2.146; 4.559

DIS, n (%)	DTD	44 (10.1)	34 (18.4)	0.0171†	0.0081	3.177	1.351; 7.469	
	Fractures and trauma	44 (10.1)	11 (5.9)		0.9565	1.028	0.384; 2.748	
	Renal and urological	37 (8.5)	9 (4.9)		-	ref	ref	
	Vascular	75 (17.2)	30 (16.2)		0.2473	1.644	0.708; 3.819	
	Infections	12 (2.8)	7 (3.8)		0.1473	2.398	0.735; 7.828	
	HN neoplasms	36 (8.3)	12 (6.5)		0.5280	1.370	0.515; 3.646	
	GI neoplasms	58 (13.3)	34 (18.4)		0.0407	2.410	1.038; 5.597	
	Other diseases	78 (17.9)	22 (11.9)		0.7384	1.160	0.486; 2.764	
	Other neoplasms	52 (11.9)	26 (14.1)		0.1035	2.056	0.863; 4.893	
RWL, n (%)	No	318 (76.4)	128 (74.9)	0.6823†	0.6823	ref	ref	
	Yes	98 (23.6)	43 (25.1)			1.090	0.721; 1.648	
SGA, n (%)	WN	343 (78.7)	131 (70.8)	0.0351†	0.0358	ref	ref	
	Malnourished	93 (21.3)	54 (29.2)			1.520	1.028; 2.248	
NRS, n (%)	At risk	165 (38.2)	81 (44.0)	0.1765†	0.1769	1.273	0.897; 1.806	
	No risk	267 (61.8)	103 (56.0)			ref	ref	
AC, n (%)	≤ P15	131 (30.3)	73 (39.9)	0.0535†	0.1936	1.359	0.856; 2.159	
	P15-P85	201 (46.5)	69 (37.7)			0.4442	0.837	0.531; 1.320
	> P85	100 (23.1)	41 (22.4)			ref	ref	
TSF, n (%)	≤ P15	65 (15.4)	33 (18.1)	0.4408†	0.2324	1.381	0.813; 2.345	
	P15-P85	220 (52.3)	99 (54.4)			0.3240	1.224	0.819; 1.829
	> P85	136 (32.3)	50 (27.5)			ref	ref	
AMC, n (%)	≤ P15	145 (34.4)	76 (41.8)	0.2306†	0.2967	1.345	0.771; 2.345	
	P15-P85	217 (51.5)	83 (45.6)			0.9454	0.981	0.569; 1.691
	> P85	59 (14.0)	23 (12.6)			ref	ref	
BMI	Overweight	186 (42.8)	85 (46.2)	0.7142†	0.5239	1.130	0.776; 1.644	
	Normal weight	178 (40.9)	72 (39.1)			-	ref	ref
	Low weight	71 (16.3)	27 (14.7)			0.8163	0.940	0.558; 1.583

Table IIA. Descriptive analysis of the variables studied and factors associated or not with weight loss during hospitalization (weight loss of at least 1 kg during hospitalization)

*Mann-Whitney test; † χ^2 -square test; ‡Univariate logistic regression. ref = reference category; OR = odds ratio; 95 % CI: OR 95 % confidence interval; HT: hospitalization time; FT: fasting time; DIS: disease; DTD: digestive tract diseases; HN: head and neck; GI: gastrointestinal; RWL: recent weight loss; SGA: subjective global assessment; WN: well nourished; NRS: nutritional risk screening; AC: arm circumference; TSF: triceps skin fold; AMC: arm muscle circumference; BMI: body mass index.

Variables	Category	No weight loss n = 436	With weight loss n = 185	p-value	p-value [†]	OR	95 % CI
%HEI/EER, n (%)	< 75 %	292 (68.7)	117 (65.4)	0.4223 [†]	0.4225	ref	0.804; 1.684
	≥ 75	133 (31.3)	62 (34.6)			1.163	
Prescribed diet, n (%)	Soft	95 (21.9)	49 (26.5)	0.1299 [†]	0.0840	1.483	0.948; 2.318
	Enteral/ Parenteral	22 (5.1)	15 (8.1)		0.0651	1.960	
	General	184 (42.5)	64 (34.6)		-	ref	0.959; 4.009
	Watery/ Fasting	107 (24.7)	41 (22.2)		0.6793	1.102	
	Liquid	25 (5.8)	16 (8.6)		0.0828	1.840	
Supplement use	No	411 (94.3)	172 (93.0)	0.5386 [†]	0.5393	ref	0.402; 1.610
	Yes	25 (5.7)	13 (7.0)			0.805	
Diet acceptance	Good	276 (79.3)	123 (79.9)	0.9893 [†]	0.8905	ref	0.553; 1.673
	Regular	49 (14.1)	21 (13.6)		0.9501	0.962	
	Poor	23 (6.6)	10 (6.5)		0.976	0.451; 2.112	
Diet change	No	228 (52.7)	88 (47.6)	0.2465 [†]	0.2468	ref	0.868; 1.731
	Yes	205 (47.3)	97 (52.4)			1.226	

Table IIB. Descriptive analysis of the variables studied and factors associated or not with weight loss during hospitalization (weight loss of at least 1 kg during hospitalization).

* χ -square test; [†]Univariate logistic regression. ref = reference category; OR: odds ratio; 95 % CI: OR 95 % confidence interval; %HEI/EER: percentage of habitual energy consumption in relation to estimated energy requirements.

DISCUSSION

This investigation showed that patients who were hospitalized for a period longer than 7 days had a three-fold greater risk of experiencing weight loss. Those who were malnourished according to the SGA had a 1.5 times greater risk. Patients diagnosed with digestive tract diseases were 3.2 times more likely to experience weight loss when compared to those with kidney and urological diseases. And patients with digestive neoplasms had a 2.4 times greater risk of presenting weight loss when compared to patients with renal and urological diseases.

A few recent studies have shown data similar to the findings of our study, such as the study by Takaoka et al. in 2017 (26); they investigated nutritional status during hospitalization using nutritional tracking instruments and laboratory exams. The authors showed that increase in hospital stay was associated with several nutritional indicators such as SGA, NRS-2002, serum albumin, and weight loss, among others (26). Another study performed by Orlandoni et al. in 2017 (27) pointed out that malnutrition was an independent predictor of length of hospital stay and mortality, and that malnourished patients were hospitalized

for almost three more days, compared with non-malnourished patients. The authors also showed that risk of death during hospitalization was 55 % higher in malnourished patients (27). Different findings were observed in the study by Cano-Torres et al., also in 2017 (28), where nutritional intervention and dietary advice contributed to a reduction in length of hospital stay but not in mortality among hospitalized malnourished patients. Other findings report that hospitalized patients may suffer from inadequate nutritional therapy, and the risk of developing malnutrition may increase during hospitalization (29).

A prospective study (30) that evaluated changes in nutritional status during hospitalization in Canada investigated SGA and body weight measurements during hospital stay and at discharge. Using multivariate analysis models, the authors showed that the decline in nutritional status according to SGA and weight loss $\geq 5\%$ were significantly associated with a longer hospital stay (30).

In an observational study conducted by Rattray et al. in 2017 (31), the nutritional adequacy of the diets prescribed and consumed by inpatients was assessed, and both supply and intake were considered adequate when they met $\geq 75\%$ of their estimated needs. In the study in question (31), it was observed that the average amount of energy and protein supplied to and consumed by the patients was significantly lower than the estimated average needs. The authors also observed that patients on liquid oral diets had a lower energy intake, showing that inpatients on liquid oral diets could develop a higher risk of malnutrition (31).

In our study, no association was observed between the type of diet that was prescribed on admission and the weight loss that occurred during hospital stay ($p = 0.1299$) (Fig. 1). Thus, we can suggest that further investigations should be carried out of hospital diets in order to better meet the energy and nutritional needs of hospitalized patients, contributing to a reduction in weight loss and hospital malnutrition. In a recent retrospective study (32) investigating the relationship between weight changes, diet, and dietary intake changes in patients with cancer, the authors found.

Variables	Category	GD n = 248	SD n = 144	LD n = 41	WDF n = 148	EPD n = 37	p-value
Age	X ± DP	56.2 ± 15.6	61.9 ± 14.9	53.6 ± 16.9	58.1 ± 14.8	56.8 ± 11.6	0.0057*§
Gender, n (%)	F	100 (40.3)	54 (37.5)	11 (26.8)	55 (37.2)	5 (13.5)	0.0193†
	M	148 (59.7)	90 (62.5)	30 (73.2)	93 (62.8)	32 (86.5)	
Disease, n (%)	DTD	9 (3.6)	15 (10.4)	12 (29.3)	39 (26.4)	2 (5.4)	< 0.0001†
	Fractures and trauma	29 (11.7)	13 (9.0)	2 (4.9)	7 (4.7)	4 (10.8)	
	Renal and urological	33 (13.3)	5 (3.5)	1 (2.4)	7 (4.7)	0 (0.0)	
	Vascular	58 (23.4)	29 (20.1)	2 (4.9)	15 (10.1)	1 (2.7)	
	Infections	10 (4.0)	4 (2.8)	0 (0.0)	3 (2.0)	2 (5.4)	
	HN neoplasms	11 (4.4)	11 (7.6)	3 (7.3)	11 (7.4)	11 (29.7)	
	DT neoplasms	9 (3.6)	24 (16.7)	11 (26.8)	33 (22.3)	15 (40.5)	
	Other diseases	51 (20.6)	24 (16.7)	5 (12.2)	18 (12.2)	1 (2.7)	
	Other neoplasms	38 (15.3)	19 (13.2)	5 (12.2)	15 (10.1)	1 (2.7)	
RWL, n (%)	No	187 (79.6)	109 (82.0)	20 (55.6)	108 (75.0)	19 (52.8)	0.0001†
	Yes	48 (20.4)	24 (18.0)	16 (44.4)	36 (25.0)	17 (47.2)	
SGA, n (%)	WN	211 (85.1)	114 (79.2)	28 (68.3)	105 (70.9)	13 (35.1)	<.0001†
	M	37 (14.9)	30 (20.8)	13 (31.7)	43 (29.1)	24 (64.9)	
NRS, n (%)	At risk	64 (26.2)	66 (45.8)	21 (51.2)	66 (44.9)	27 (73.0)	<.0001†
	No risk	180 (73.8)	78 (54.2)	20 (48.8)	81 (55.1)	10 (27.0)	
AC, n (%)	≤ P15	72 (29.4)	38 (26.4)	16 (39.0)	52 (35.9)	25 (67.6)	0.0002†
	P15-P85	119 (48.6)	64 (44.4)	15 (36.6)	60 (41.4)	11 (29.7)	
	> P85	54 (22.0)	42 (29.2)	10 (24.4)	33 (22.8)	1 (2.7)	
TSF, n (%)	≤ P15	32 (13.2)	26 (18.3)	7 (17.1)	19 (13.9)	14 (37.8)	0.0125†
	P15-P85	132 (54.3)	69 (48.6)	18 (43.9)	80 (58.4)	17 (45.9)	
	>P85	79 (32.5)	47 (33.1)	16 (39.0)	38 (27.7)	6 (16.2)	
AMC, n (%)	≤ P15	78 (32.0)	44 (31.0)	20 (48.8)	53 (39.0)	25 (67.6)	0.0003†
	P15-P85	132 (54.1)	81 (57.0)	16 (39.0)	58 (42.6)	12 (32.4)	
	> P85	34 (13.9)	17 (12.0)	5 (12.2)	25 (18.4)	0 (0.0)	
BMI	Overweight	126 (51.2)	61 (42.4)	17 (41.5)	61 (41.2)	5 (13.5)	0.0022†
	Normal weight	87 (35.4)	62 (43.1)	18 (43.9)	62 (41.9)	19 (51.4)	
	Low weight	33 (13.4)	21 (14.6)	6 (14.6)	25 (16.9)	13 (35.1)	

Table III. Comparison of study variables by types of prescribed diet,

*Kruskal-Wallis test; †χ-square test; ‡Fisher's exact test; §Differences between soft and general diets; soft and liquid. GD: general diet; SD: soft diet; LD: liquid diet; WDF: watery diet and fasting; EPD: enteral and parenteral diet; F: female; M: male; DTD: digestive tract disease; HN: head and neck; DT: digestive tract; Dis: diseases; RWL: recent weight loss; SGA: subjective global assessment; WN: well nourished; M: malnourished; NRS: nutritional risk screening; AC: arm circumference; TSF: triceps skin fold; AMC: arm muscle circumference; BMI: body mass index.

A significant positive correlation between dietary energy or protein consumption and weight change. Another finding of the study was a correlation between anorexia and weight loss; but the authors found no similar correlation between changes in dietary intake and changes in weight (32). The study also pointed out that patients with advanced cancer and weight loss could be consuming diets insufficient to maintain body weight (32). In any case, it is essential to monitor dietary consumption and the evolution of body weight throughout hospitalization. A prospective observational study (33) that investigated the clinical characteristics and long-term results of unintentional weight loss in outpatients, showed the relevance of monitoring long-term clinical evolution in patients with unexplained weight loss (33).

And in a prospective study in a Brazilian hospital (34), factors related to the reduction of prescribed enteral therapy were investigated, showing that operational problems, gastric stasis, and accidental loss of the gastric tube, among others, were the main reasons for inadequate food intake.




Figure 1. Percentage distribution of the type of diet prescribed on admission, according to weight loss (loss or no loss of at least 1 kg weight), during hospital stay: $p = 0.1299$ (χ -square).

The findings of the present study are interesting and can easily illustrate clinical scenarios for several underlying conditions. Our findings also point out the importance of investigating weight loss, of assessing type of diet and food consumption, and of daily, continuous monitoring of the nutritional aspects of hospitalized patients. Some works are in line with the proposal of our investigation, such as a cross-sectional study (35) to investigate the tracking of malnutrition through anthropometric and laboratory indicators in patients admitted to medical wards. The authors observed a longer hospital stay in

malnourished patients, evidencing that malnutrition was more common among patients with malignancies (35).

It is important to note that the variables and/or all nutritional indicators investigated in the present study are currently instruments in use for the nutritional assessment and monitoring of hospitalized patients, and have been used in other investigations reported in the relevant literature. This is the case of a study that assessed the prevalence of malnutrition (29) and determined the daily caloric intake of hospitalized patients. The nutritional risk found by the NRS was 44.6 % (29), an index similar to that found in the present investigation, where we verified a 39.9 % rate of nutritional risk. As for malnutrition by SGA, the authors found a 53.6 % rate of malnutrition (29) while in our study it was 23.6 %. Possibly, the malnutrition found here by the SGA may have been underestimated in relation to the nutritional risk obtained by the NRS.

Finally, in the present study it was observed that all the variables studied showed a significant association with the type of diet prescribed at hospital admission (Table III). Although no direct relationship was found between diet and weight loss during hospital stay (Table IIB), due to the fact that we found these differences in relation to the type of diet prescribed, we can suggest that “indirectly” there is an association of weight loss during hospitalization with the prescribed diet. The diet prescribed was not directly related to weight loss but through the factors that are associated with diet prescription (in these factors there was a significant difference).

CONCLUSION

Malnutrition, diseases, digestive neoplasms, and length of stay were associated with weight loss during hospitalization. The findings of this study highlight the relevance of nutritional and dietary monitoring of hospitalized patients.

STUDY LIMITATIONS

This study had some limitations inherent to the type of retrospective study, in relation to other prospective and controlled studies, with loss of some nutritional variable data. That is why there was less information for some of the nutritional variables analyzed when compared to others. On the other hand, retrospective studies have the advantage that all the nutritional parameters that were evaluated reflected the actual conditions of routine hospital care, as it actually happens in daily hospital clinical practice, and not in a controlled manner, which could make the outcome artificial.

ACKNOWLEDGMENTS

We are grateful for the support received from PUC-Campinas (Campinas, SP, Brazil).

FUNDING

This project was not sponsored.

CONFLICTS OF INTEREST

The authors have no conflicts of interest to declare.

ETHICAL APPROVAL FOR RETROSPECTIVE STUDIES

This project was approved by the institution's ethics committee (reference number: 2.312.714; Caae: 75798017.0.0000.5481).

REFERENCES

1. Correia MITD, Perman MI, Waitzberg DL. Hospital malnutrition in Latin America: A systematic review. *Clin Nutr* 2017;36:958-67. DOI: 10.1016/j.clnu.2016.06.025
2. Khalatbari-Soltani S, Marques-Vidal P. Impact of nutritional risk screening in hospitalized patients on management, outcome and costs: A retrospective study. *Clin Nutr* 2016;35(6):1340-6. DOI: 10.1016/j.clnu.2016.02.012
3. Silva FR, de Oliveira MG, Souza AS, Figueroa JN, Santos CS. Factors associated with malnutrition in hospitalized cancer patients: a cross-sectional study. *Nutr J* 2015; 10;14:123. DOI: 10.1186/s12937-015-0113-1
4. Cruz V, Bernal L, Buitrago G, Ruiz AJ. Screening for malnutrition among hospitalized patients in a Colombian University Hospital. *Rev Med Chil* 2017;145(4):449-57. DOI: 10.4067/S0034-98872017000400005
5. Luma HN, Eloumou SAFB, Mboligong FN, Temfack E, Donfack OT, Doualla MS. Malnutrition in patients admitted to the medical wards of the Douala General Hospital: a cross-sectional study. *BMC Res Notes* 2017;10(1):238. DOI: 10.1186/s13104-017-2592-y
6. Cederholm T, Bosaeus I, Barazzoni R, Bauer J, Van Gossum A, Klek S, et al. Diagnostic criteria for malnutrition. An ESPEN Consensus Statement. *Clin Nutr* 2015;34:335-40. DOI: 10.1016/j.clnu.2015.03.001
7. Bonilla-Palomas JL, Gámez-López AL, Castillo-Domínguez JC, Moreno-Conde M, López Ibáñez MC, Alhambra Expósito R, et al. Nutritional Intervention in Malnourished Hospitalized Patients with Heart Failure. *Arch Med Res* 2016;47(7):535-40. DOI: 10.1016/j.arcmed.2016.11.005
8. O'Shea E, Trawley S, Manning E, Barrett A, Browne V, Timmons S. Malnutrition in Hospitalised Older Adults: A Multicentre Observational Study of Prevalence, Associations and Outcomes. *J Nutr Health Aging* 2017;21(7):830-6. DOI: 10.1007/s12603-016-0831-x

9. Kiss N. Nutrition support and dietary interventions for patients with lung cancer: current insights. *Lung Cancer (Auckl)* 2016;7:1-9. DOI: 10.2147/LCTT.S85347
10. Zheng H, Huang Y, Shi Y, Chen W, Yu J, Wang X. Nutrition status, nutrition support therapy, and food intake are related to prolonged hospital stays in China: Results from the Nutrition Day 2015 Survey. *Ann Nutr Metab* 2016;69(3-4):215-25. DOI: 10.1159/000451063
11. Mortensen MN, Larsen AK, Skadhaug LB, Høgsted RH, Beermann T, Cook ME, et al. Protein and energy intake improved by in-between meals: An intervention study in hospitalized patients. *Clin Nutr ESPEN* 2019;30:113-8. DOI: 10.1016/j.clnesp.2019.01.007
12. Hedman S, Nydahl M, Faxén-Irving G. Individually prescribed diet is fundamental to optimize nutritional treatment in geriatric patients. *Clin Nutr* 2016;35(3):692-8. DOI: 10.1016/j.clnu.2015.04.018
13. Detsky AS, McLaughlin JR, Baker JP, Johnston N, Whittaker S, Mendelson RA, et al. What is subjective global assessment of nutritional status? *JPEN* 1987;11:8-13. DOI: 10.1177/014860718701100108
14. Kondrup J, Allison SP, Elia M, Vellas B, Plauth M. ESPEN guidelines for nutrition screening 2002. *Clin Nutr* 2003;22(4):415-21. DOI: 10.1016/S0261-5614(03)00098-0
15. Kondrup J, Rasmussen HH, Hamberg O, Stanga Z, ESPEN Working Group. Nutritional risk screening (NRS 2002): a new method based on a analysis of controlled clinical trials. *Clin Nutr* 2003;22(3):321-36. DOI: 10.1016/S0261-5614(02)00214-5
16. World Health Organization (WHO). Obesity: Preventing and managing the global epidemic - Report of a WHO Consultation on obesity. Geneva: WHO; 1998.
17. Organização Pan-Americana de Saúde. XXXVI Reunião del Comitê Asesor de Investigaciones en Salud – Encuesta Multicêntrica – Salud Beinestar y Envejecimeiento (SABE) en América Latina e el Caribe. Informe preliminar, Washington, D.C. OPAS; 2001. Available from: <http://www.opas.org/program/sabe.htm>.
18. Frisancho AR. Anthropometric standards for the assessment of growth and nutritional status. Michigan: The University of Michigan Press; 1990. DOI: 10.3998/mpub.12198
19. Burr ML, Phillips MK. Anthropometric norms in the elderly. *Br J Nutr* 1984;51:165-9. DOI: 10.1079/BJN19840020
20. Harris J, Benedict F. A biometric study of basal metabolism in man. Washington D.C. Carnegie Institute of Washington; 1919.
21. Sorensen J, Kondrup J, Prokopowicz J, Schiesser M, Krahenbuhl L, Méier R, et al. EuroOOPS: An international, multicentre study to implement nutritional risk screening and evaluate clinical outcome. *Clin Nutr* 2008;27:340-9. DOI: 10.1016/j.clnu.2008.03.012
22. Miriran P, Hosseinpour-Niazi S, Mehrabani HH, Kavian F, Azizi F. Validity and reliability of a nutrition screening tool in hospitalized patients. *Nutrition* 2011;27(6):647-52. DOI: 10.1016/j.nut.2010.06.013
23. Conover WJ. Practical Nonparametric Statistics. John Wiley & Sons Inc. Nova Iorque; 1971.

24. Tabachnick BG, Fidell LS. Using Multivariate Statistics. Boston: Allyn and Bacon, 4th ed; 2001, pp. 966.
25. SAS System for Windows (Statistical Analysis System), versão 9.4. SAS Institute Inc, 2002-2012, Cary, NC, USA.
26. Takaoka A, Sasaki M, Nakanishi N, Kurihara M, Ohi A, Bamba S, et al. Nutritional Screening and Clinical Outcome in Hospitalized Patients with Crohn's Disease. *Ann Nutr Metab* 2017;71(3-4):266-72. DOI: 10.1159/000485637
27. Orlandoni P, Venturini C, Jukic Peladic N, Costantini A, Di Rosa M, Cola C, et al. Malnutrition upon Hospital Admission in Geriatric Patients: Why Assess It? *Front Nutr* 2017;30:4:50. DOI: 10.3389/fnut.2017.00050
28. Cano-Torres EA, Simental-Mendía LE, Morales-Garza LA, Ramos-Delgado JM, Reyes-Gonzalez MM, Sánchez-Nava VM, et al. Impact of Nutritional Intervention on Length of Hospital Stay and Mortality among Hospitalized Patients with Malnutrition: A Clinical Randomized Controlled Trial. *J Am Coll Nutr* 2017;36(4):235-9. DOI: 10.1080/07315724.2016.1259595
29. Konturek PC, Herrmann HJ, Schink K, Neurath MF, Zopf Y. Malnutrition in Hospitals: It Was, Is Now, and Must Not Remain a Problem! *Med Sci Monit* 2015;21:2969-75. DOI: 10.12659/MSM.894238
30. Allard JP, Keller H, Jeejeebhoy KN, Laporte M, Duerksen DR, Gramlich L, et al. Decline in nutritional status is associated with prolonged length of stay in hospitalized patients admitted for 7 days or more: A prospective cohort study. *Clin Nutr* 2016;35(1):144-52. DOI: 10.1016/j.clnu.2015.01.009
31. Rattray M, Desbrow B, Roberts S. Comparing nutritional requirements, provision and intakes among patients prescribed therapeutic diets in hospital: An observational study. *Nutrition* 2017;39:40:50-56. DOI: 10.1016/j.nut.2017.03.006
32. Nasrah R, Kanbalian M, Van Der Borch C, Swinton N, Wing S, Jagoe RT. Defining the role of dietary intake in determining weight change in patients with cancer cachexia. *Clin Nutr* 2018;37(1):235-41. DOI: 10.1016/j.clnu.2016.12.012
33. Bosch X, Monclús E, Escoda O, Guerra-García M, Moreno P, Guasch N, et al. Unintentional weight loss: Clinical characteristics and outcomes in a prospective cohort of 2677 patients. *PLoS One* 2017;12(4):e0175125. DOI: 10.1371/journal.pone.0175125
34. Martins JR, Shiroma GM, Horie LM, Logullo L, Silva M de L, Waitzberg DL. Factors leading to discrepancies between prescription and intake of enteral nutrition therapy in hospitalized patients. *Nutrition* 2012;28(9):864-7. DOI: 10.1016/j.nut.2011.07.025
35. Luma HN, Eloumou SAFB, Mboligong FN, Temfack E, Donfack OT, Doualla MS. Malnutrition in patients admitted to the medical wards of the Douala General Hospital: a cross-sectional study. *BMC Res Notes* 2017;10(1):238. DOI: 10.1186/s13104-017-2592-y

ÍNDICE REMISSIVO

A

Ácidos graxos 109, 110, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 127, 128, 168, 169, 171, 172, 286

Adesão 7, 10, 51, 52, 53, 54, 56, 58, 59, 79, 82, 84, 205, 207, 214, 246

Adolescência 92, 211, 241, 242, 248, 250, 254

Alta hospitalar 10, 223, 226

Assistência domiciliar 217, 219, 224, 226, 227, 228, 230

Assistência hospitalar 223, 268

Atenção básica à saúde 108, 255, 264, 265

Atenção farmacêutica 25, 26, 32, 33, 35, 38, 39, 40, 41, 43, 46, 47, 48

Atuação do farmacêutico 1, 3, 7, 26, 31, 50

Audição 162, 163, 168, 169, 172, 173

Automedicação 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 29, 30, 31, 32, 34, 35, 36, 37, 41, 49, 263

Avaliação nutricional 192, 194, 196, 197, 198

C

Camellia sinensis 153, 154, 156, 157, 159, 160

Canabidiol 130, 131, 134, 135, 136

Cannabis 130, 131, 133, 134, 135

CBD 130, 131, 133, 134

Centro cirúrgico 15, 266, 268, 269, 270, 276, 277, 278

Centro de Atenção Psicossocial 210, 212, 213

Chá verde 153, 154, 155, 156, 157, 158, 159, 160, 161

Consumo de medicamentos 3, 11, 25, 26, 29, 36, 38, 40, 41, 44, 47, 48

Contraceptivo de emergência 86, 87, 88, 90, 93

Controle 2, 9, 25, 26, 27, 28, 36, 40, 41, 44, 45, 54, 59, 98, 100, 126, 132, 159, 162, 164, 165, 227, 237, 246, 247, 248, 250, 262, 266, 280, 281

Cuidado 7, 9, 11, 33, 49, 73, 74, 84, 90, 132, 178, 179, 180, 181, 182, 183, 184, 187, 188, 189, 190, 200, 202, 203, 204, 205, 208, 210, 211, 212, 213, 214, 215, 216, 218, 219, 220, 223, 224, 225, 226, 227, 228, 229, 230, 231, 261, 267, 276, 277, 278

Cuidados farmacêuticos 61

D

Diabetes mellitus 6, 66, 192, 193, 194, 196, 198, 199, 200, 201, 241, 242, 243, 244, 252, 253

Dieta 40, 62, 63, 64, 69, 72, 73, 75, 113, 114, 115, 125, 127, 137, 138, 157, 158, 161, 162,

163, 164, 168, 171, 174, 194, 197, 198, 199, 247, 248, 280, 284, 286
Disbiose 279, 280, 281, 282, 283, 284, 285, 286, 287, 288
Distúrbios endócrinos 241, 243, 251
Doença de alzheimer 217, 218, 219, 222

E

Educação à distância 233
Educação em saúde 13, 84, 220, 221, 228, 231, 234, 238, 262, 263
Efeitos adversos 1, 7, 8, 25, 30, 34, 58, 63, 78, 79, 86, 134, 155
Enfermagem 3, 12, 13, 14, 15, 16, 17, 20, 21, 22, 23, 36, 49, 72, 73, 74, 75, 76, 77, 78, 79, 80, 82, 83, 84, 85, 91, 92, 94, 127, 207, 208, 209, 220, 221, 222, 230, 231, 232, 235, 254, 258, 264, 265, 266, 268, 269, 270, 271, 272, 274, 276, 277, 278, 287
Epidemiologia 11, 22, 49, 192, 200, 209, 254
Epilepsia 130, 131, 132, 133, 134, 135, 136
Equipe interdisciplinar de saúde 61, 213
Equipe multiprofissional 71, 73, 204, 205, 207, 212, 223, 272, 278
Eventos adversos 8, 64, 178, 179, 180, 182, 183, 188, 189, 202, 203, 204, 206, 208, 209

F

Fitoterápicos 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108

H

HIV 52, 53, 54, 55, 56, 57, 58, 59, 60, 89, 179

I

Idoso fragilizado 217, 219
Idosos 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 44, 49, 192, 193, 194, 195, 196, 197, 198, 199, 200, 220
Infância 130, 132, 211, 241, 242, 243, 246, 248, 250, 253, 254
Inquéritos 280

L

Legislação 34, 81, 83, 95, 98, 99, 103, 104, 105, 108, 215
Lipídios 109, 110, 111, 113, 118, 119, 122, 124

M

Medicamentos antirretrovirais 52, 53, 54, 55, 56

O

Obesidade 153, 154, 155, 156, 157, 158, 159, 160, 161, 163, 193, 194, 195, 197, 199, 220, 241, 242, 244, 245, 246, 247, 248, 249, 250, 251, 252, 281, 283, 285, 286, 287, 288

P

Palmeiras 109, 110, 111, 123, 129
Pediatria 136, 241, 251, 252, 254
Perda auditiva 162, 163, 164, 167, 169, 171, 172, 173, 174, 250
Perfil de medicamentos 25
Pílula do dia seguinte 86, 87, 93
Plantas medicinais 95, 96, 97, 98, 100, 101, 102, 103, 104, 105, 106, 107, 108, 155
Políticas de saúde 23, 95, 96
Prevenção 10, 14, 25, 33, 36, 59, 64, 80, 81, 82, 83, 102, 123, 128, 154, 155, 157, 158, 159, 162, 163, 164, 168, 171, 172, 192, 211, 213, 219, 227, 228, 241, 246, 250, 251, 254, 263, 280
Prevenção de doenças 33, 158, 192, 228, 280
Primeiros socorros 232, 233, 234, 235, 236, 237, 238, 239, 240
Psicotrópicos 38, 39, 40, 41, 42, 43, 44, 46, 47, 48, 49

Q

Qualidade da assistência à saúde 178, 207
Qualidade em saúde 179, 181, 182, 188, 203
Questionários 255, 257, 258, 259, 263, 279, 280, 282, 283, 284, 285, 287

R

Relações comunidade-instituição 233

S

Saúde Mental 41, 50, 210, 211, 212, 213, 214, 215, 216
Saúde Ocupacional 78, 83, 255
Segurança do paciente 9, 36, 61, 74, 178, 179, 180, 182, 183, 185, 187, 188, 190, 203, 204, 205, 206, 207, 208, 209, 268, 277
Serviços de saúde 16, 17, 42, 79, 178, 179, 181, 182, 184, 188, 196, 202, 203, 204, 205, 210, 211, 252
Stress 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 277

U

Uso de medicamentos 4, 5, 6, 9, 10, 13, 14, 18, 21, 23, 25, 26, 27, 29, 30, 32, 49, 54, 61, 64, 73, 74, 131, 133
Uso descontrolado 86, 87
Uso racional de medicamentos 1, 11, 25, 26, 30, 31, 32, 34, 35, 37, 38, 39, 41

V

Vias de administração de medicamentos 61
Visita domiciliar 223, 224, 225, 226, 228, 229, 230, 231.

 www.arenaeditora.com.br
 contato@arenaeditora.com.br
 [@arenaeditora](https://www.instagram.com/arenaeditora)
 www.facebook.com/arenaeditora.com.br

Ciências da saúde:

Políticas públicas, assistência e gestão


Ano 2021

 www.atenaeditora.com.br
 contato@atenaeditora.com.br
 [@atenaeditora](https://www.instagram.com/atenaeditora)
 www.facebook.com/atenaeditora.com.br

Ciências da saúde:

Políticas públicas, assistência e gestão