

CIÊNCIAS DA SAÚDE:

Oferta, acesso e utilização 2




Edson da Silva
Rodrigo Lellis Santos
(Organizadores)

Atena
Editora
Ano 2022

CIÊNCIAS DA SAÚDE:

Oferta, acesso e utilização 2



Edson da Silva
Rodrigo Lellis Santos
(Organizadores)

Atena
Editora
Ano 2022

Editora chefe

Profª Drª Antonella Carvalho de Oliveira

Editora executiva

Natalia Oliveira

Assistente editorial

Flávia Roberta Barão

Bibliotecária

Janaina Ramos

Projeto gráfico

Camila Alves de Cremo

Daphynny Pamplona

Gabriel Motomu Teshima

Luiza Alves Batista

Natália Sandrini de Azevedo

Imagens da capa

iStock

Edição de arte

Luiza Alves Batista

2022 by Atena Editora

Copyright © Atena Editora

Copyright do texto © 2022 Os autores

Copyright da edição © 2022 Atena Editora

Direitos para esta edição cedidos à Atena Editora pelos autores.

Open access publication by Atena Editora



Todo o conteúdo deste livro está licenciado sob uma Licença de Atribuição *Creative Commons*. Atribuição-Não-Comercial-NãoDerivativos 4.0 Internacional (CC BY-NC-ND 4.0).

O conteúdo dos artigos e seus dados em sua forma, correção e confiabilidade são de responsabilidade exclusiva dos autores, inclusive não representam necessariamente a posição oficial da Atena Editora. Permitido o *download* da obra e o compartilhamento desde que sejam atribuídos créditos aos autores, mas sem a possibilidade de alterá-la de nenhuma forma ou utilizá-la para fins comerciais.

Todos os manuscritos foram previamente submetidos à avaliação cega pelos pares, membros do Conselho Editorial desta Editora, tendo sido aprovados para a publicação com base em critérios de neutralidade e imparcialidade acadêmica.

A Atena Editora é comprometida em garantir a integridade editorial em todas as etapas do processo de publicação, evitando plágio, dados ou resultados fraudulentos e impedindo que interesses financeiros comprometam os padrões éticos da publicação. Situações suspeitas de má conduta científica serão investigadas sob o mais alto padrão de rigor acadêmico e ético.

Conselho Editorial**Ciências Biológicas e da Saúde**

Profª Drª Aline Silva da Fonte Santa Rosa de Oliveira – Hospital Federal de Bonsucesso

Profª Drª Ana Beatriz Duarte Vieira – Universidade de Brasília

Profª Drª Ana Paula Peron – Universidade Tecnológica Federal do Paraná

Prof. Dr. André Ribeiro da Silva – Universidade de Brasília

Profª Drª Anelise Levay Murari – Universidade Federal de Pelotas

Prof. Dr. Benedito Rodrigues da Silva Neto – Universidade Federal de Goiás



Prof. Dr. Cirêno de Almeida Barbosa – Universidade Federal de Ouro Preto
Prof^o Dr^a Daniela Reis Joaquim de Freitas – Universidade Federal do Piauí
Prof^o Dr^a Débora Luana Ribeiro Pessoa – Universidade Federal do Maranhão
Prof. Dr. Douglas Siqueira de Almeida Chaves – Universidade Federal Rural do Rio de Janeiro
Prof. Dr. Edson da Silva – Universidade Federal dos Vales do Jequitinhonha e Mucuri
Prof^o Dr^a Elizabeth Cordeiro Fernandes – Faculdade Integrada Medicina
Prof^o Dr^a Eleuza Rodrigues Machado – Faculdade Anhanguera de Brasília
Prof^o Dr^a Elane Schwinden Prudêncio – Universidade Federal de Santa Catarina
Prof^o Dr^a Eysler Gonçalves Maia Brasil – Universidade da Integração Internacional da Lusofonia Afro-Brasileira
Prof. Dr. Ferlando Lima Santos – Universidade Federal do Recôncavo da Bahia
Prof^o Dr^a Fernanda Miguel de Andrade – Universidade Federal de Pernambuco
Prof. Dr. Fernando Mendes – Instituto Politécnico de Coimbra – Escola Superior de Saúde de Coimbra
Prof^o Dr^a Gabriela Vieira do Amaral – Universidade de Vassouras
Prof. Dr. Gianfábio Pimentel Franco – Universidade Federal de Santa Maria
Prof. Dr. Helio Franklin Rodrigues de Almeida – Universidade Federal de Rondônia
Prof^o Dr^a Iara Lúcia Tescarollo – Universidade São Francisco
Prof. Dr. Igor Luiz Vieira de Lima Santos – Universidade Federal de Campina Grande
Prof. Dr. Jefferson Thiago Souza – Universidade Estadual do Ceará
Prof. Dr. Jesus Rodrigues Lemos – Universidade Federal do Piauí
Prof. Dr. Jônatas de França Barros – Universidade Federal do Rio Grande do Norte
Prof. Dr. José Aderval Aragão – Universidade Federal de Sergipe
Prof. Dr. José Max Barbosa de Oliveira Junior – Universidade Federal do Oeste do Pará
Prof^o Dr^a Juliana Santana de Curcio – Universidade Federal de Goiás
Prof^o Dr^a Lívia do Carmo Silva – Universidade Federal de Goiás
Prof. Dr. Luís Paulo Souza e Souza – Universidade Federal do Amazonas
Prof^o Dr^a Magnólia de Araújo Campos – Universidade Federal de Campina Grande
Prof. Dr. Marcus Fernando da Silva Praxedes – Universidade Federal do Recôncavo da Bahia
Prof^o Dr^a Maria Tatiane Gonçalves Sá – Universidade do Estado do Pará
Prof. Dr. Maurilio Antonio Varavallo – Universidade Federal do Tocantins
Prof^o Dr^a Mylena Andréa Oliveira Torres – Universidade Ceuma
Prof^o Dr^a Natiéli Piovesan – Instituto Federaci do Rio Grande do Norte
Prof. Dr. Paulo Inada – Universidade Estadual de Maringá
Prof. Dr. Rafael Henrique Silva – Hospital Universitário da Universidade Federal da Grande Dourados
Prof^o Dr^a Regiane Luz Carvalho – Centro Universitário das Faculdades Associadas de Ensino
Prof^o Dr^a Renata Mendes de Freitas – Universidade Federal de Juiz de Fora
Prof^o Dr^a Sheyla Mara Silva de Oliveira – Universidade do Estado do Pará
Prof^o Dr^a Suely Lopes de Azevedo – Universidade Federal Fluminense
Prof^o Dr^a Vanessa da Fontoura Custódio Monteiro – Universidade do Vale do Sapucaí
Prof^o Dr^a Vanessa Lima Gonçalves – Universidade Estadual de Ponta Grossa
Prof^o Dr^a Vanessa Bordin Viera – Universidade Federal de Campina Grande
Prof^o Dr^a Welma Emidio da Silva – Universidade Federal Rural de Pernambuco



Ciências da saúde: oferta, acesso e utilização 2

Diagramação: Camila Alves de Cremo
Correção: Yaidy Paola Martinez
Indexação: Amanda Kelly da Costa Veiga
Revisão: Os autores
Organizadores: Edson da Silva
Rodrigo Lellis Santos

Dados Internacionais de Catalogação na Publicação (CIP)

C569 Ciências da saúde: oferta, acesso e utilização 2 /
Organizadores Edson da Silva, Rodrigo Lellis Santos. -
Ponta Grossa - PR: Atena, 2022.

Formato: PDF

Requisitos de sistema: Adobe Acrobat Reader

Modo de acesso: World Wide Web

Inclui bibliografia

ISBN 978-65-258-0052-3

DOI: <https://doi.org/10.22533/at.ed.523222303>

1. Ciências da saúde. I. Silva, Edson da (Organizador).
II. Santos, Rodrigo Lellis (Organizador). III. Título.

CDD 613

Elaborado por Bibliotecária Janaina Ramos – CRB-8/9166

Atena Editora
Ponta Grossa – Paraná – Brasil
Telefone: +55 (42) 3323-5493
www.atenaeditora.com.br
contato@atenaeditora.com.br



Atena
Editora
Ano 2022

DECLARAÇÃO DOS AUTORES

Os autores desta obra: 1. Atestam não possuir qualquer interesse comercial que constitua um conflito de interesses em relação ao artigo científico publicado; 2. Declaram que participaram ativamente da construção dos respectivos manuscritos, preferencialmente na: a) Concepção do estudo, e/ou aquisição de dados, e/ou análise e interpretação de dados; b) Elaboração do artigo ou revisão com vistas a tornar o material intelectualmente relevante; c) Aprovação final do manuscrito para submissão.; 3. Certificam que os artigos científicos publicados estão completamente isentos de dados e/ou resultados fraudulentos; 4. Confirmam a citação e a referência correta de todos os dados e de interpretações de dados de outras pesquisas; 5. Reconhecem terem informado todas as fontes de financiamento recebidas para a consecução da pesquisa; 6. Autorizam a edição da obra, que incluem os registros de ficha catalográfica, ISBN, DOI e demais indexadores, projeto visual e criação de capa, diagramação de miolo, assim como lançamento e divulgação da mesma conforme critérios da Atena Editora.



DECLARAÇÃO DA EDITORA

A Atena Editora declara, para os devidos fins de direito, que: 1. A presente publicação constitui apenas transferência temporária dos direitos autorais, direito sobre a publicação, inclusive não constitui responsabilidade solidária na criação dos manuscritos publicados, nos termos previstos na Lei sobre direitos autorais (Lei 9610/98), no art. 184 do Código Penal e no art. 927 do Código Civil; 2. Autoriza e incentiva os autores a assinarem contratos com repositórios institucionais, com fins exclusivos de divulgação da obra, desde que com o devido reconhecimento de autoria e edição e sem qualquer finalidade comercial; 3. Todos os e-book são *open access*, *desta forma* não os comercializa em seu site, sites parceiros, plataformas de *e-commerce*, ou qualquer outro meio virtual ou físico, portanto, está isenta de repasses de direitos autorais aos autores; 4. Todos os membros do conselho editorial são doutores e vinculados a instituições de ensino superior públicas, conforme recomendação da CAPES para obtenção do Qualis livro; 5. Não cede, comercializa ou autoriza a utilização dos nomes e e-mails dos autores, bem como nenhum outro dado dos mesmos, para qualquer finalidade que não o escopo da divulgação desta obra.



APRESENTAÇÃO

A coletânea '*Ciências da saúde: oferta, acesso e utilização*' é uma obra composta por 44 capítulos, organizados em dois volumes. Ambos abordam diferentes áreas de conhecimento no campo da saúde. Os autores compartilham resultados de seus projetos acadêmicos ou de atuações profissionais. Além disso, alguns capítulos são ensaios teóricos ou revisões sobre a temática.

A coletânea conta com as contribuições de discentes e docentes de vários cursos de graduação e de pós-graduação, bem como outros profissionais de instituições que estabeleceram parcerias com as universidades envolvidas.

O volume 2 reúne 24 capítulos com autoria multidisciplinar. Nota-se a importância da atuação interdisciplinar, revelando os avanços nesse campo do ensino superior no Brasil. As vivências compartilhadas corroboram com a consolidação das atividades acadêmicas que integram, cada vez mais, universidades, instituições e as comunidades envolvidas.

Esperamos que as vivências relatadas nessa obra contribuam para o enriquecimento da formação universitária e da atuação profissional com o fortalecimento das práticas interdisciplinares nas ciências da saúde. Agradecemos aos autores que tornaram essa coletânea possível e lhe desejamos uma ótima leitura.

Edson da Silva
Rodrigo Lellis Santos

SUMÁRIO

CAPÍTULO 1..... 1

YOUTUBE™ COMO FONTE DE INFORMAÇÕES SOBRE DIABETES: É TUDO FAKE NEWS?

Edson da Silva

Rodrigo Lellis Santos

Ana Luísa Simões Guedes

 <https://doi.org/10.22533/at.ed.5232223031>

CAPÍTULO 2..... 9

PROFISSIONAIS E ACADÊMICOS DO EIXO SAÚDE – O ENTENDIMENTO DA AUTO MEDICAÇÃO COMO RISCO À SAÚDE PESSOAL: REVISÃO BIBLIOGRÁFICA

Rosecley Santana Bispo

Thatielle Baldez de Oliveira

Ethienny Baldez de Oliveira Pacheco

Gabriel Rodrigues dos Santos

Rodrigo Lima dos Santos Pereira

Viviane Pires do Nascimento

João Marcos Torres do Nascimento Mendes

Axell Donelli Leopoldino Lima


Paula Lauane Araújo

Sueli Pereira de Sousa

Brenda Soares Coêlho

Isabela Carvalho Tupy

Lustarllone Bento de Oliveira

 <https://doi.org/10.22533/at.ed.5232223032>

CAPÍTULO 3..... 25

A PREVALÊNCIA DOS ESTUDOS SOBRE ESPIRITUALIDADE NA ÁREA DA SAÚDE

Ivando Amancio da Silva Junior

Adelaide Souza da Silva Rodrigues

Eronildo de Andrade Braga

Jânio Marcio de Sousa

José Ednésio Cruz Freire

Lucimar Camelo Souza Silva

Madna Avelino Silva


Romildo Alves Batista

Samuel Ramalho Torres Maia

Givanildo Carneiro Benício

Germana Maria Viana Cruz

Ticiano Maria Lima Azevedo

 <https://doi.org/10.22533/at.ed.5232223033>

CAPÍTULO 4..... 35

PSICOSE PUERPERAL

Danielle Freire Goncalves

Carlito dias da Silva
José Wneyldson da Silveira
Isaac Prado Ramos
Iara Priscilla Inácio de Freitas
Mariana Hoover Miranda Rezende
Gabriela Cordeiro Silva
Sarah da Silva Barros
José Danilo Amorim Ghidetti
Paloma de Faria Guerra
Thiago Mourão Almeida Araújo
Francimar Neto de Almeida Lopes

 <https://doi.org/10.22533/at.ed.5232223034>

CAPÍTULO 5..... 41

MANEJO DO PÉ DIABÉTICO NA ATENÇÃO BÁSICA

Luiza Schinke Genn

 <https://doi.org/10.22533/at.ed.5232223035>

CAPÍTULO 6..... 53

A QUALIDADE DE VIDA E O ENFRENTAMENTO DA DOENÇA DE MULHERES COM DIAGNÓSTICO DE CÂNCER DE MAMA


Guilherme Vinício de Sousa Silva
Angela Makeli Kososki Dalagnol
Keroli Eloiza Tessaro da Silva
Débora Tavares de Resende e Silva

 <https://doi.org/10.22533/at.ed.5232223036>

CAPÍTULO 7..... 59

PRINCIPAIS TÉCNICAS MOLECULARES UTILIZADAS PARA VERIFICAR A COMPATIBILIDADE HLA ENTRE DOADOR E RECEPTOR NO TRANSPLANTE DE RINS PROVENIENTES DE DOADOR FALECIDO: UMA REVISÃO


Camilla Natália Oliveira Santos
Lucas Sousa Magalhães

 <https://doi.org/10.22533/at.ed.5232223037>

CAPÍTULO 8..... 72

A ASSISTÊNCIA EM PACIENTES COM CÂNCER DE MAMA NO SISTEMA ÚNICO DE SAÚDE – SUS


Jacqueline Aragão de Medeiros

 <https://doi.org/10.22533/at.ed.5232223038>

CAPÍTULO 9..... 80

FATORES DE RISCO, CAUSAS, MANIFESTAÇÕES DA GAGUEIRA INFANTIL: UMA REVISÃO SISTEMÁTICA


Isadora Cássia de Oliveira
Mariana Ferraz Conti Uvo

 <https://doi.org/10.22533/at.ed.5232223039>

CAPÍTULO 10..... 98

ASSOCIAÇÃO ENTRE INFECÇÃO E COINFECÇÃO POR PAPILOMAVÍRUS HUMANO (HPV) E EPSTEIN-BARR VÍRUS (EBV) E CÂNCERES DE CAVIDADE ORAL, OROFARINGE E NASOFARINGE

Pietriny Emanuelli Piana
Vítor Nakayam Shiguemoto
Rosebel Trindade Cunha Prates
Léia Carolina Lucio

 <https://doi.org/10.22533/at.ed.52322230310>

CAPÍTULO 11 103

PROPOSTA PEDAGÓGICA PARA A EDUCAÇÃO FÍSICA, NA FORMA HÍBRIDA, NA EDUCAÇÃO BÁSICA


Marcus Tullius de Paula Senna
Carlos Roberto Alves Teles

 <https://doi.org/10.22533/at.ed.52322230311>

CAPÍTULO 12..... 116

INFLUÊNCIAS DA ACREDITAÇÃO INTERNACIONAL NO ENFRENTAMENTO DA COVID-19 EM UM HOSPITAL PRIVADO DE BELO HORIZONTE: RELATO DE EXPERIÊNCIA


Camila Martins de Jesus
Stéphane Bruna Barbosa
Karla Rona da Silva
Fátima Ferreira Roquete

 <https://doi.org/10.22533/at.ed.52322230312>

CAPÍTULO 13..... 127

CONTRIBUIÇÕES DA PERMANÊNCIA DO ACOMPANHANTE A PACIENTES INTERNADOS EM UNIDADE DE TERAPIA INTENSIVA (UTI) PEDIÁTRICA: RELATO DE EXPERIÊNCIA

Gisele da Silva Peixoto Zandoná
Camila Fortes Correa
Nádia Dan Bianchi de Souza
Patrick Jean Barbosa Sales
Ana Carolini Ferreira de Castro
Shanna Machado de Sousa
Lucia Helaynn Penha de Souza Franco


 <https://doi.org/10.22533/at.ed.52322230313>

CAPÍTULO 14..... 137

RELATO DE CASO: NÓDULO MAMÁRIO NA PARACOCCIDIOIDOMICOSE

Carina Pereira Bigheti
Eduardo Carvalho Pessoa
Paulo Eduardo Hernandes Antunes
Suzana Shinomia
Paulo Henrique Pedroso de Lima

Lucas Golçalves Cardoso
Leandro Clementino Falcão
Ana Laura Lopes Potente
Erika Mayumi Watanabe
Maria Célia Franco Issa
Gabriela Ferreira Bailão
Murilo Bucci Vega

 <https://doi.org/10.22533/at.ed.52322230314>

CAPÍTULO 15..... 150

CORPO LÍQUIDO: PROBLEMATIZAÇÕES SOBRE CIRURGIAS ESTÉTICAS NA MODERNIDADE E AVALIAÇÕES PSICOLÓGICAS

Everley Rosane Goetz
Carolina Guidi Gentil

 <https://doi.org/10.22533/at.ed.52322230315>

CAPÍTULO 16..... 158

LEVANTAMENTO DAS GUIAS DE TRATAMENTO COM ANTIDEPRESSIVOS E ANÁLISE DOS MEDICAMENTOS DISPONIBILIZADOS PELO SUS NO MUNICÍPIO DE GUARAPUAVA-PR


Mariana Hyeda Miranda
Luana Mota Ferreira
Daniel De Paula

 <https://doi.org/10.22533/at.ed.52322230316>

CAPÍTULO 17..... 171

CARACTERÍSTICAS FÍSICO-QUÍMICAS E BIOLÓGICAS DA CELULOSE BACTERIANA DA CANA-DE-AÇÚCAR

Emerson Leonardo de Moura Santos
Veridiana Sales Barbosa de Souza
Rodrigo Pontes Lima
Anderson Arnaldo Silva
Ana Olívia de Andrade e Souza
Carlos Eduardo de Souza Rodrigues
Adriana Parente Vianna Simões Ferreira
Kristian Pires Gurgel
Márcio Handerson Benevides de Freitas
Mariana Cavalcanti Pirajá Viana Ferreira
Olávio Campos Júnior
Amanda Vasconcelos de Albuquerque

 <https://doi.org/10.22533/at.ed.52322230317>

CAPÍTULO 18..... 185

AVALIAÇÃO DOS BENEFÍCIOS E DA SEGURANÇA DA UTILIZAÇÃO COSMÉTICA DO ÓLEO DE COCO *IN NATURA* PARA PELE E CABELO

Jackeline de Souza Alecrim
Mariane Parma Ferreira de Souza

Tathiana Gomes Chaves


 <https://doi.org/10.22533/at.ed.52322230318>

CAPÍTULO 19.....200

ASSÉDIO MORAL NAS RELAÇÕES DE TRABALHO: IDENTIFICAÇÃO E ANÁLISE DAS CONSEQUÊNCIAS PARA A SAÚDE DOS SERVIDORES

Mirely Ferreira dos Santos

Livia Maria Duarte de Castro

 <https://doi.org/10.22533/at.ed.52322230319>

CAPÍTULO 20.....213

CARACTERÍSTICAS CLÍNICAS DA DOENÇA DE VON WILLEBRAND: UMA REVISÃO BIBLIOGRÁFICA INTEGRATIVA


Lydia Gabriela Fooshang Bustillos

Diego Brito Dos Santos

Fernanda Letícia Rodrigues

Juan Pereira da Silva

Rayssa Gabrielle Pereira de Castro Bueno

 <https://doi.org/10.22533/at.ed.52322230320>

CAPÍTULO 21.....221

EXERCÍCIOS DE VIBRAÇÃO DE CORPO INTEIRO COMO INTERVENÇÃO PARA ADULTOS SOBREVIVENTES DE CÂNCER: REVISÃO SISTEMÁTICA

Ana Gabriellie Valério Penha

Dayana Figueiredo Genovez da Silva

Ester Fonseca de Melo

Fabiana Jóia da Silva Nunes

Luelia Teles Jaques de Albuquerque


Ana Carolina Coelho-Oliveira

Juliana Pessanha de Freitas

Márcia Cristina Moura-Fernandes

Mario Bernardo-Filho

Danúbia da Cunha de Sá-Caputo

 <https://doi.org/10.22533/at.ed.52322230321>

CAPÍTULO 22.....238

ANÁLISIS DE LA INFLUENCIA DE LA ANSIEDAD EN FUNCIÓN DEL GÉNERO Y LA EDAD EN DEPORTISTAS DE DOMA CLÁSICA

María Merino Fernández

Michelle Matos Duarte

Rafael Alarcón Guerrero

Pilar Jerez Villanueva

Bárbara Rodríguez Rodríguez


 <https://doi.org/10.22533/at.ed.52322230322>

CAPÍTULO 23.....251

ROUX-EN-Y GASTRIC BYPASS IMPROVES IN SHORT TERM THE CLINICAL-

**ANTHROPOMETRIC PARAMETERS AND REDUCES RISK FOR OBESITY-RELATED
CARDIOMETABOLIC DISEASES**


Thiago da Rosa Lima
Paula Caroline de Almeida
Fabrício Azevedo Voltarelli
Lilian Culturato
Eudes Thiago Pereira Ávila
Wender Junior de Deus Silva
James Wilfred Navalta
Amilcar Sabino Damazo

 <https://doi.org/10.22533/at.ed.52322230323>

CAPÍTULO 24..... 263

EWINGS SARCOMA THE ILIAC BONE - REPORT OF CASE

Ricardo Dias Borges
Emanuella Chaves De Moura

 <https://doi.org/10.22533/at.ed.52322230324>

SOBRE OS ORGANIZADORES 271

ÍNDICE REMISSIVO..... 272

ROUX-EN-Y GASTRIC BYPASS IMPROVES IN SHORT TERM THE CLINICAL-ANTHROPOMETRIC PARAMETERS AND REDUCES RISK FOR OBESITY-RELATED CARDIOMETABOLIC DISEASES

Data de aceite: 01/03/2022

Amilcar Sabino Damazo

Federal University of Mato Grosso. Cuiabá. Mato Grosso. Brazil; Graduation Program in Health Sciences. Faculty of Medicine. Brazil

Thiago da Rosa Lima

Federal University of Mato Grosso. Cuiabá. Mato Grosso. Brazil; Graduation Program in Health Sciences. Faculty of Medicine. Brazil University Center of Várzea Grande. Várzea Grande. Mato Grosso, Brazil
<https://orcid.org/0000-0001-8544-6797>

Paula Caroline de Almeida

Federal University of Mato Grosso. Cuiabá. Mato Grosso. Brazil; Graduation Program in Health Sciences. Faculty of Medicine. Brazil

Fabrcio Azevedo Voltarelli

Federal University of Mato Grosso. Cuiabá. Mato Grosso. Brazil; Graduation Program in Health Sciences. Faculty of Medicine. Brazil

Lilian Culturato

University Center of Várzea Grande. Várzea Grande. Mato Grosso, Brazil

Eudes Thiago Pereira Ávila

Federal University of Mato Grosso. Cuiabá. Mato Grosso. Brazil; Graduation Program in Health Sciences. Faculty of Medicine. Brazil

Wender Junior de Deus Silva

University Center of Várzea Grande. Várzea Grande. Mato Grosso, Brazil

James Wilfred Navalta

University of Nevada-Las Vegas
Las Vegas, Nevada, USA

ABSTRACT: BACKGROUND: Roux-en-Y gastric bypass surgery (RYGB) is the most applied technique in the treatment of severe obesity worldwide. However, its impact on anthropometric parameters and the risk for cardiometabolic diseases in obese patients is uncertain.

OBJECTIVE: To evaluate anthropometric clinical parameters and the evolution of risk factors for obesity-related diseases in individuals of both sexes undergoing RYGB. **METHODOLOGY:**

Sixty-nine adults subjects from both sexes submitted to RYGB surgery treatment were divided into 3 groups: G1(<13 months, n=24); G2 (>13 and <25 months, n=21), and G3 (>25 and <37 months, n=24). Sociodemographic and anthropometric information before and after surgery were collected. The abdominal perimeter was used in the classification of cardiometabolic risk and the BMI was used for the risk of obesity-related diseases. Hypotheses were tested by Student's t-test and ANOVA, and the significance level adopted was 5%. **RESULTS:** The average age was 36.0±10.0 years, with 69.6% being male and 30.4% female. Anthropometric parameters (weight, BMI, and abdominal circumference) were higher among women, except for weight loss and percentage of weight loss. There was a difference in weight loss between the sexes in the moments before and after RYGB. There was a decrease in the risk of disease due to

obesity and cardiovascular diseases after RYGB. Weight loss and %WL were greater years by year in the short term of 3 years after surgery. **CONCLUSION:** RYGB proved to be an effective strategy for both sexes in combating obesity, providing in the short term a significant improvement in clinical-anthropometric parameters and reduction of risk factors for obesity-related cardiometabolic diseases.

KEYWORDS: Bariatric Surgery. Severe obesity. Nutritional and metabolic diseases.

BYPASS GÁSTRICO ROUX-EN-Y MELHORA A CURTO PRAZO OS PARÂMETROS CLÍNICO-ANTROPOMÉTRICOS E REDUZ O RISCO DE DOENÇAS CARDIOMETABÓLICAS RELACIONADAS À OBESIDADE

RESUMO: INTRODUÇÃO: cirurgia de bypass gástrico em Y-de-Roux (BGYR) é a técnica mais aplicada no tratamento da obesidade grave em todo o mundo. No entanto, seu impacto sobre os parâmetros antropométricos e o risco de doenças cardiometabólicas em pacientes obesos é incerto. **OBJETIVO:** Avaliar os parâmetros clínicos antropométricos e a evolução dos fatores de risco para doenças relacionadas à obesidade em indivíduos de ambos os sexos submetidos ao BGYR. **METODOLOGIA:** Sessenta e nove indivíduos adultos de ambos os sexos submetidos ao tratamento cirúrgico com BGYR foram divididos em 3 grupos: G1 (<13 meses, n = 24); G2 (> 13 e <25 meses, n = 21) e G3 (> 25 e <37 meses, n = 24). Foram coletadas informações sociodemográficas e antropométricas antes e após a cirurgia. O perímetro abdominal foi utilizado para a classificação do risco cardiometabólico e o IMC para o risco de doenças relacionadas à obesidade. As hipóteses foram testadas pelo teste t de Student e ANOVA, e o nível de significância adotado foi de 5%. **RESULTADOS:** A média de idade foi de 36,0±10,0 anos, sendo 69,6% do sexo masculino e 30,4% do feminino. Os parâmetros antropométricos (peso, IMC e circunferência abdominal) foram maiores entre as mulheres, exceto para perda de peso e %PP. Houve diferença na perda de peso entre os sexos nos momentos antes e depois do BGYR. Houve diminuição do risco de doenças por obesidade e doenças cardiovasculares após o BGYR. A perda de peso e %PP foram maiores ano a ano no curto prazo (3 anos após a cirurgia). **CONCLUSÃO:** O BGYR demonstrou ser uma estratégia eficaz para ambos os sexos no combate à obesidade, proporcionando em curto prazo uma melhora significativa dos parâmetros clínico-antropométricos e redução dos fatores de risco para doenças cardiometabólicas relacionadas à obesidade.

PALAVRAS-CHAVE: Cirurgia bariátrica. Obesidade severa. Doenças metabólicas e nutricionais.

INTRODUCTION

The increasing prevalence of obesity at all stages of life has a serious impact on public health worldwide (1). Besides, not just excess body mass by subcutaneous and intra-abdominal fat (visceral) accumulation but countless other diseases are triggered in association with obesity such as diabetes mellitus, cardiovascular diseases, osteoporosis, and cancer (2).

Body mass index (BMI) is a clinical tool widely used in the identification of obesity (2). Individuals with a BMI of 25.0 to 29.9 kg/m² are considered overweight; obese when they

have a BMI of 30 kg/m² or higher being BMI of 40 kg/m² or more is considered as severe obesity (3). It is widely recognized that the increase in BMI is directly associated with the onset of chronic diseases and mortality from various causes (4). It is estimated that with each increase of 5 kg/m² from the eutrophic state (i.e., 25 kg/m²) there is an increase of 29% in general mortality 41% in vascular mortality, and 210% in mortality related to diabetes (5).

Epidemiological data indicate that in low-income countries obesity is prevalent among middle-aged adults. whereas in high-income countries this predominance occurs among individuals of both sexes and varied ages (6). In Brazil, on account of being a developing country (i.e., low income) recent data collected in the Surveillance of Risk and Protection Factors for Chronic Diseases by Telephone Survey (VIGITEL) revealed a higher incidence of obesity in young adults aged between 25 and 34 years old (84.2%) and middle-aged adults between 35 and 44 years old (81.1%). When considering sex, women with 20.7% stand out showing a higher prevalence of obesity than men with 18.7% (7).

Guidelines for the treatment of obesity in several countries, including Brazil, recommend multidisciplinary clinical treatment through behavioral therapy, regular physical exercise, and low-calorie diets to mitigate negative psychological effects reduce body fat, increase lean mass. in addition to attenuating the comorbidities generated by excess fat composing to the treatment, if necessary, the use of specific drugs (8-10).

However, the vast majority of cases of severe obesity do not present sufficient outcomes with conventional clinical treatment. requiring the adoption of surgical alternatives (11). In this context. the Roux-en-Y gastric by-pass (RYGB) technique is considered one of the most efficient methods and incurs a lower morbidity and mortality rate (12, 13). Among the various bariatric and metabolic surgery techniques, RYGB has been considered the gold standard is the most popular and the most adopted considered a well-tolerated procedure in the control of obesity (14).

Based on the above, the present study aimed to evaluate clinical parameters and the evolution of risk factors for obesity-related diseases in individuals of both sexes who underwent Roux-en-Y gastric bypass surgery in the city of Cuiabá/MT between January 2015 and January 2018.

MATERIALS AND METHODS

This is a cross-sectional study through a retrospective analysis of the database of patients undergoing RYGB surgery. The study was previously approved by the Research Ethics Committee of the Academic Center of Várzea Grande-MT (CEP/Univag) with protocol n° 2.959.596.

The study consisted of data collection from 69 adult patients, of both sexes, aged between 18 and 65 years who were followed up in a particular establishment specialized in the clinical and surgical treatment of obesity, metabolic and digestive diseases located in

the county of Cuiabá/MT throughout January 2015 and January 2018.

The inclusion criteria adopted were bariatric and metabolic surgery using the RYGB technique, and the premise of having undergone regular multi-professional clinical follow-up to assess surgical viability was adopted in addition to maintaining the follow-up for at least 3 years after surgery.

The information collected through medical records, medical history, anamnesis, anthropometric measurements, and nutritional assessment were: sex, age, professional occupation, education level, race, presence or absence of comorbidities, the time elapsed after surgery, body weight before of surgery (Usual weight - UW), bodyweight after surgery (current weight - CW) as well as the usual and current abdominal perimeters.

The level of education was classified according to that proposed by the Ministry of Education (15). Race /color was self-reported and followed the proposal in the color and race classification system described by the Brazilian Institute of Geography and Statistics (IBGE) (16). For the evaluation of metabolic and cardiovascular risk, obtained due to the excessive accumulation of visceral fat, the measurement of the abdominal perimeter (waist region) was adopted and classified as proposed by the WHO (3). The body mass index (BMI) obtained using the formula , was used to evaluate the risk of diseases associated with obesity (i.e.. comorbidities caused by excess weight and body fat) were classified as proposed by the Brazilian Obesity Guidelines (2016) (8). The percentage of body weight loss (%WL) was obtained using the formula (17). Regarding the time elapsed after surgery the participants were divided into 3 groups: G1 (<13 months; n=24); G2 (> 13 and <25 months; n=21), and G3 (> 25 and <37 months; n=24) for comparisons between weight loss (kg) % WL and BMI.

Descriptive analysis of qualitative data was expressed as a percentage, and the associations between these variables the χ^2 -square test were used. Quantitative variables were analyzed for normality using the Kolmogorov-Smirnov method. Student's t-test for independent samples was used for analysis between genders. Parameters such as body weight, % WL and BMI were analyzed by Student's t-test for dependent samples (before and after surgery). Data regarding the time elapsed after surgery were tested by analysis of variance (one-way ANOVA) and when necessary Tukey's post-hoc was applied. The results are expressed as mean \pm standard deviation (SD), and the significance level adopted was 5%. All analyzes were performed using software the Statistical Package for Social Sciences (SPSS®) version 25.

RESULTS

Sociodemographic profile

The data regarding the sociodemographic profile are presented in table 1. In a

total of 69 participants with a mean age (36.0 ± 10.0 years) previously submitted to RYGB surgery, it was observed that 48 (69.6 %) were male and 21 female (30.4%). Regarding race 37 participants reported being brown (53.6%), 22 white (31.9%), 8 participants (11.6%) considered themselves black, and 2 yellow (2.9%). Education level was 4 (5.8%) participants with incomplete elementary school, 8 (11.6%) complete elementary school, 25 (36.2%) complete high school, and 32 individuals (46.4%) with complete undergraduate degree. The professional occupation of the participants portrayed that 42 of them (60.9%) were employed, 17 (24.6%) were unemployed, 9 were self-employed (13.0%) and only 1 (1.4%) was retired.

Sex	Frequency	Percentage
	n	%
Male	48	69.6
Female	21	30.4
Race/color	n	%
White	22	31.9
Black	8	11.6
Yellow	2	2.9
Brown	37	53.6
Education levels	n	%
Incomplete elementary school	4	5.8
Complete elementary school	8	11.6
Complete high school	25	36.2
Complete undergraduate degree	32	46.4
Professional occupation	n	%
Employed	42	60.9
Unemployed	17	24.6
Self-employed	9	13.0
Retired	1	1.4

Table 1 Sociodemographic profile of the studied population.
Data presented as relative frequency (n) and percentage (%).

Anthropometric clinical parameters

Anthropometric clinical parameters are shown in table 2. The usual weight ($p < 0.001$) and the current weight ($p < 0.001$) were higher in females. However, there was no difference for the usual BMI between genders ($p = 0.06$). The current BMI was lower for males ($p = 0.03$). There was no difference in the usual abdominal perimeter ($p = 0.06$). However, the current abdominal perimeter was lower among males ($p = 0.02$). Weight loss (kg) ($p = 0.11$) and weight loss percentage ($p = 0.67$) were similar between genders.

Variables		$\bar{X}\pm DP$	Minimum	Maximum	<i>p</i> -value
Usual weight (kg ⁻¹)	Male	105.6±12.1	83.0	140.0	<0.001
	Female	127.1±19.7	89.0	168.0	
Current weight (kg ⁻¹)	Male	75.2±13.6	53.0	121.0	<0.001
	Female	91.3±13.7	70.0	118.0	
Usual BMI (kg/m ²)	Male	39.8±4.2	33.1	50.8	0.06
	Female	42.0±5.2	34.7	54.2	
Current BMI (kg/m ²)	Male	28.1±4.3	20.9	41.9	0.03
	Female	30.6±4.3	24.5	39.0	
Usual abdominal perimeter (cm)	Male	117.0±9.7	94.0	136.0	0.06
	Female	124.0±5.2	115.0	131.0	
Current abdominal perimeter (cm)	Male	92.4±13.6	73.0	120.0	0.02
	Female	101.0±9.9	77.0	113.0	
Weight loss (kg)	Male	30.4±11.6	6.0	49.0	0.11
	Female	35.8±15.3	9.0	67.0	
Weight loss (%)	Male	28.7±10.2	7.2	43.7	0.67
	Female	27.5±9.6	7.3	41.6	

Table 2 Anthropometric clinical parameters

Student t-test for independent samples and a 5% significance level. Data presented as mean±standard deviation ($\bar{X}\pm DP$).

Analysis of anthropometric parameters in the before and after RYGB

There was a reduction between before and after surgery moments of RYGB for both men and women on anthropometric parameters of body weight (differences of mean: men = -30.4 kg; women = -35.8 kg) (figure 1A); BMI (differences of mean: men = -11.7 kg/m²; women = -11.4 kg/m²) (figure 1B); and abdominal perimeter (differences of mean: men = -24.3 cm; women = -22.4 cm) (figure 1C).




Figure 1 Anthropometric parameters in the before and after RYGB moments. The data were evaluated by Student's t-test for dependent (paired) samples, and a significance level of 5% was adopted. The results are presented as mean ± standard deviation ($\bar{X}\pm DP$). The differences between the means are presented as the mean of the post-surgery moment - mean of the pre-surgical moment (B - A).

Association of risks for obesity-related diseases and the time elapsed from RYGB

The association of risks for diseases caused by excess body weight and accumulation of abdominal fat before surgery and between the after-surgery period is shown in figure 3. There was an association between the risk of diseases due to obesity before surgery ($p=0.02$). In a total of 69 participants, it was observed that 7 (10.1%) were at high risk, 28 (40.6%) at very high risk, and 34 (49.3%) at extremely high risk for comorbidities associated with obesity. There was no association between the risk of diseases associated with obesity and the time after surgery ($p=0.23$). Similarly, there was no association between cardiovascular risk in the before surgery period ($p=0.31$). There was an association between cardiovascular risk and after surgery time ($p=0.01$), where, in a total of 60 participants, 23 (38.3%) had normal risk, 6 (10.0%) high risk, and 31 (51.7%) extremely high risk for cardiovascular diseases.

Comorbidities	Classification	Groups (%)			Total (%)	χ^2 <i>p-value</i>
		G1	G2	G3		
Risk of disease due to before surgery obesity	High	5 (71.4)	2 (28.6)	0 (0.0)	7 (10.1)	0.02
	Very high	13 (46.4)	8 (28.6)	7 (25.0)	28 (40.6)	
	Extremely high	6 (17.6)	11(32.4)	17 (50.0)	34 (49.3)	
Risk of disease due to after surgery obesity	Normal	2 (15.4)	3 (23.1)	8 (61.5)	13 (18.8)	0.23
	Little high	9 (29.0)	11 (35.5)	11 (35.5)	31 (44.9)	
	High	9 (50.0)	6 (33.3)	3 (16.7)	18 (26.1)	
	Very high	3 (50.0)	1 (16.7)	2 (33.3)	6 (8.7)	
CD risk before surgery	Extremely high	1 (100)	0 (0.0)	0 (0.0)	1 (1.4)	0.31
	Very high	0 (0.0)	2 (66.7)	1 (33.3)	3 (4.9)	
	Extremely high	20 (34.5)	17 (29.3)	21(36.2)	58 (95.1)	
CD risk after surgery	Normal	13 (56.5)	6 (26.1)	4 (17.4)	23 (38.3)	0.01
	High	1 (16.7)	4 (66.7)	1 (16.7)	6 (10.0)	
	Very high	6 (19.4)	10 (32.3)	15 (48.4)	31 (51.7)	

Table 3 Association of risk for diseases caused by excess body weight and cardiovascular diseases (CVD) due to the accumulation of abdominal fat in the before and after RYGB.

Groups indicating the time elapsed post-surgery G1 (<13 months. n = 24). G2 (> 13 and <25 months. n=21). G3 (> 25 and <37 months. n=24). The classification of risk factors (comorbidities) was performed according to the Brazilian Obesity Guidelines (2016) [18]. The data were submitted to association analysis by chi-square (χ^2) with a significance level of 5%. The results are presented as relative frequency (n) and Percentage (%)

Total weight loss and percentage of weight loss concerning the elapsed time of RYGB

Weight loss (kg) was higher in G3 (44.4 ± 7.0) compared to G1 (19.0 ± 8.3) and G2 (32.8 ± 7.0) as well as G2 was higher than G1 ($p<0.0001$) (figure 2A). The percentage of weight loss was higher in the G3 group (36.5 ± 5.4) compared to G1 and G2 (19.1 ± 9.2 ; 29.5 ± 4.8), respectively; and the G2 was greater than G1 ($p<0.0001$) (figure 2B).




Figure 2 (A) Total weight loss (kg); and (B) percentage of weight loss. Groups indicating the time elapsed after surgery G1 (<13 months, n=24), G2 (> 13 and <25 months; n=21), G3 (> 25 and <37 months, n=24). The data were evaluated by analysis of variance (Anova one-way) and when necessary Tukey HSD's post-hoc presented in superscript letters, was applied significance level of 5%.

DISCUSSION

The present study aimed to evaluate anthropometric clinical parameters and the evolution of risk factors for obesity-related diseases in adult individuals of both sexes who underwent bariatric and metabolic surgery using the RYGB technique. The main findings show that RYGB induced, in both sexes, an improvement in the nutritional profile and a reduction in risk factors for cardiovascular and metabolic diseases associated with obesity. The study portrayed a greater number of male patients who underwent bariatric and metabolic surgery for RYGB, which at least for this technique contradicts the overall frequency of predominance of bariatric and metabolic surgery by women as reported in the literature (19).

Notably, the positive outcomes observed were similar between men and women undergoing RYGB bariatric and metabolic surgery when we consider the classic anthropometric parameters expected after the procedure such as rapid weight loss, reduced BMI and morbidities related to the degree of obesity, reduction cardiovascular risk is seen by the smaller abdominal perimeter and an increase in the %WL (19-21). Although widely studied, there is still no consensus on which metrics should be adopted to confirm a good or bad response to short and long-term weight loss and recovery. Even so, the surgical treatment of severe obesity and super obesity is considered effective, especially when it provides sustained reductions in excess body weight and BMI (11).

The purposes of bariatric and metabolic surgery, especially RYGB, are based on the improvement of clinical parameters such as weight loss, reduced risk of morbidities, remission and/or control of metabolic diseases, and improved quality of life (22, 23). In general, the sustained loss of excess body weight is understood as one of the main parameters to determine the success and/or failure of a strategy, surgical or not, in the treatment of obesity, and this outcome is considered essential to improve clinical conditions of the individual (21, 24). At this point, even though the present study evaluated patients

with a maximum time of up to 3 years after RYGB, the observed results indicate significant clinical benefits, at least concerning anthropometric clinical parameters.

Among the different techniques of surgical approach, RYGB is considered the gold standard in the treatment of severe obesity and super obesity, since its outcome provides a rapid and considerable reduction in excess body weight, which favors the control of metabolic morbidities, attenuating the risk of deleterious cardiovascular events and overlapping perioperative risk (25). In the present study, we observed a positive change in the classification of risk factors for diseases caused by excess body weight and for cardiovascular diseases (CD) due to the accumulation of abdominal fat concerning the time of surgery, which corroborates the recent findings in the literature (26-28).

The risk of developing catastrophic cardiovascular events is greatly increased according to the degree of obesity (4). An interesting study sought to evaluate the cardiovascular risk estimated in 10 years of obese patients considering the moments before and 2 years after the treatment of obesity with RYGB. The study involved 42 patients; of these, the majority were female, 85.7% with a mean age of 43.5 ± 8.6 years. The results registered a significant reduction in cardiovascular risk associated, mainly, with the reduction of body weight and improvement of obesity-related morbidities such as diabetes mellitus and arterial hypertension. According to the authors, super-obese patients benefited from the effects of RYGB surgery, especially over the first 2 years that followed, this outcome is related to the accelerated weight loss that occurs in this period (29).

Throughout the treatment of severe obesity, rapid weight loss becomes a key measure in preventing metabolic and cardiovascular risk in the short term, and long term if sustained. There are reports in the literature that in the immediate post-surgical period of RYGB, patients have excellent responses on clinical parameters (14, 30). In a clinical trial conducted with 4047 obese individuals, divided between those undergoing bariatric and metabolic surgery and those who were treated conventionally, the results showed that after 2 years, there was an increase in body weight by 0.1% for the control group and a reduction of 23.4% for the surgical group. However, after 10 years, weight increased by 1.6% for the control and decreased by 16.1% for the surgery group. In summary, the group undergoing surgery had lower incidence rates for diabetes, dyslipidemia, and hyperuricemia after 2 and 10 years (28). However, it is still contradictory to say that the benefits remain over time or whether the disease will tend to return, and the serious risks for morbidities may still reappear (31).

Sociodemographic, socio-cultural and psychosocial aspects have a strong impact on eating behavior, energy intake, and physical activity level (30). Therefore, so far, there is no known single factor that can be attributed to sustained weight change, but rather a set of interdisciplinary elements that stimulate positive physiological, genetic, behavioral, and motivational responses. In this sense, we understand that special care should be implemented over time after a surgical follow-up of Roux-en-Y gastric bypass, since the

effects of the technique lead to energy restriction and malabsorption of nutrients, among other events that can cause damage to short and/or long term.

In the present study, it was evidenced that the individuals participating in the research presented satisfactory results, even though the short time after surgery is a bias factor to confirm this information. The main limitations of the study are in the cross-sectional design, which prevents the monitoring and makes it impossible to obtain new data for further analyzes even more robust. Certainly, some additional clinical data such as blood pressure, blood glucose, lipid profile, body composition could be included for a better analysis of the outcome. However, concerning the central proposal of the study, the results described manage in one way or another to contribute to the understanding of the clinical course of patients with a high degree of obesity and who underwent surgical treatment of obesity using the RYGB technique.

CONCLUSION

The data described here support the assumption that bariatric and metabolic surgery using the Roux-en-Y gastric bypass technique proved to be an effective strategy for both sexes in combating obesity, providing, in the short term, a significant improvement in the anthropometric clinical parameters as fast weight loss, increased %WL, reduced BMI and risk factors for cardiometabolic and obesity-related diseases. Future research is important to provide new knowledge aimed at assisting the clinical follow-up of obese individuals on the way to surgical treatment or those who have already been treated surgically aiming mainly at health and quality of life.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ACKNOWLEDGEMENTS

This research did not receive any specific grant from funding agencies in the public or not-for-profit sectors. The authors acknowledge CAPES, FAPEMAT, CNPq, and Academic Center of Varzea Grande/MT (UNIVAG).

REFERENCES

1. Engin A. The Definition and Prevalence of Obesity and Metabolic Syndrome. *Advances in experimental medicine and biology*. 2017;960:1-17. doi:10.1007/978-3-319-48382-5_1
2. Apovian CMJAJMC. Obesity: definition, comorbidities, causes, and burden. 2016;22(7 Suppl):s176-85.

3. WHO. Physical status: the use and interpretation of anthropometry. Report of a WHO Expert Committee. *World Health Organization technical report series*. 1995;854:1-452.
4. Global BMIMC, Di Angelantonio E, Bhupathiraju Sh N, et al. Body-mass index and all-cause mortality: individual-participant-data meta-analysis of 239 prospective studies in four continents. *Lancet (London, England)*. Aug 20 2016;388(10046):776-86. doi:10.1016/s0140-6736(16)30175-1
5. Whitlock G, Lewington S, Sherliker P, et al. Body-mass index and cause-specific mortality in 900 000 adults: collaborative analyses of 57 prospective studies. *Lancet (London, England)*. Mar 28 2009;373(9669):1083-96. doi:10.1016/s0140-6736(09)60318-4
6. Swinburn BA, Sacks G, Hall KD, et al. The global obesity pandemic: shaped by global drivers and local environments. *Lancet (London, England)*. Aug 27 2011;378(9793):804-14. doi:10.1016/s0140-6736(11)60813-1
7. Brasil. VIGITEL Brasil 2018: vigilância de fatores de risco e proteção para doenças crônicas por inquérito telefônico: estimativas sobre frequência e distribuição sociodemográfica de fatores de risco e proteção para doenças crônicas nas capitais dos 26 estados brasileiros e no Distrito Federal em 2018. In: Saúde Md, editor.: SVS/MS Brasília; 2019.
8. ABESO. Diretrizes Brasileiras de Obesidade. 4ª ed. São Paulo, SP: ABESO 2016.
9. Ryan DH, Kahan S. Guideline Recommendations for Obesity Management. *The Medical clinics of North America*. Jan 2018;102(1):49-63. doi:10.1016/j.mcna.2017.08.006
10. Yumuk V, Tsigos C, Fried M, et al. European Guidelines for Obesity Management in Adults. *Obesity facts*. 2015;8(6):402-24. doi:10.1159/000442721
11. Jastrzębska-Mierzyńska M, Ostrowska L, Hady HR, Dadan J, Konarzewska-Duchnowska E. The impact of bariatric surgery on nutritional status of patients. *Wideochirurgia i inne techniki maloinwazyjne = Videosurgery and other miniinvasive techniques*. Apr 2015;10(1):115-24. doi:10.5114/wiitm.2014.47764
12. Kahan S. Overweight and obesity management strategies. *Am J Manag Care*. Jun 2016;22(7 Suppl):s186-96.
13. Lim CH, Jahansouz C, Abraham AA, Leslie DB, Ikramuddin S. The future of the Roux-en-Y gastric bypass. *Expert review of gastroenterology & hepatology*. Jul 2016;10(7):777-84. doi:10.1586/17474124.2016.1169921
14. Navarrete S, Leyba JL, LI SN, Borjas G, Tapia JL, Alcázar R. Results of The Comparative Study of 200 Cases: One Anastomosis Gastric Bypass vs Roux-en-Y Gastric Bypass. *Obesity surgery*. Sep 2018;28(9):2597-2602. doi:10.1007/s11695-018-3224-x
15. Brasil. Lei de Diretrizes e Bases da Educação Nacional. In: Federal S, editor.: Brasília; 2005.
16. Osório RG. O sistema classificatório de cor ou raça do IBGE. 2003;
17. Blackburn GL, Harvey KB. Nutritional assessment as a routine in clinical medicine. *Postgraduate medicine*. May 1982;71(5):46-63. doi:10.1080/00325481.1982.11716062

18. Mancini M. Diretrizes Brasileiras de Obesidade 2016. *Associação Brasileira para o Estudo da Obesidade e da Síndrome Metabólica 4ª ed São Paulo*. 2016;
19. Kennedy-Dalby A, Adam S, Ammori BJ, Syed AA. Weight loss and metabolic outcomes of bariatric surgery in men versus women - A matched comparative observational cohort study. *European journal of internal medicine*. Dec 2014;25(10):922-5. doi:10.1016/j.ejim.2014.10.020
20. Sherf Dagan S, Goldensluger A, Globus I, et al. Nutritional Recommendations for Adult Bariatric Surgery Patients: Clinical Practice. *Adv Nutr*. 2017;8(2):382-394. doi:10.3945/an.116.014258
21. Wolfe BM, Kvach E, Eckel RH. Treatment of Obesity: Weight Loss and Bariatric Surgery. *Circulation research*. May 27 2016;118(11):1844-55. doi:10.1161/circresaha.116.307591
22. Brethauer SA, Kim J, el Chaar M, et al. Standardized outcomes reporting in metabolic and bariatric surgery. *Surgery for obesity and related diseases : official journal of the American Society for Bariatric Surgery*. May-Jun 2015;11(3):489-506. doi:10.1016/j.soard.2015.02.003
23. de Mattos LC, Costa ACC. Clinical and laboratory review of the reversion of metabolic syndrome in patients submitted to bariatric surgery after one year of the procedure. *Brazilian Journal of Development*. 2021;7(2):18278-95.
24. Grover BT, Morell MC, Kothari SN, Borgert AJ, Kallies KJ, Baker MT. Defining Weight Loss After Bariatric Surgery: a Call for Standardization. *Obesity surgery*. Nov 2019;29(11):3493-3499. doi:10.1007/s11695-019-04022-z
25. Buchwald H, Avidor Y, Braunwald E, et al. Bariatric surgery: a systematic review and meta-analysis. *Jama*. Oct 13 2004;292(14):1724-37. doi:10.1001/jama.292.14.1724
26. Schiavon CA, Bersch-Ferreira AC, Santucci EV, et al. Effects of Bariatric Surgery in Obese Patients With Hypertension: The GATEWAY Randomized Trial (Gastric Bypass to Treat Obese Patients With Steady Hypertension). *Circulation*. Mar 13 2018;137(11):1132-1142. doi:10.1161/circulationaha.117.032130
27. Mundbjerg LH, Stolberg CR, Cecere S, et al. Supervised Physical Training Improves Weight Loss After Roux-en-Y Gastric Bypass Surgery: A Randomized Controlled Trial. *Obesity (Silver Spring, Md)*. May 2018;26(5):828-837. doi:10.1002/oby.2214328.
28. Sjöström L, Lindroos AK, Peltonen M, et al. Lifestyle, diabetes, and cardiovascular risk factors 10 years after bariatric surgery. *The New England journal of medicine*. Dec 23 2004;351(26):2683-93. doi:10.1056/NEJMoa035622
29. Donadelli SP, Salgado W, Jr., Marchini JS, et al. Change in predicted 10-year cardiovascular risk following Roux-en-Y gastric bypass surgery: who benefits? *Obesity surgery*. May 2011;21(5):569-73. doi:10.1007/s11695-010-0348-z
30. Christensen BJ, Schmidt JB, Nielsen MS, et al. Patient profiling for success after weight loss surgery (GO Bypass study): An interdisciplinary study protocol. *Contemporary clinical trials communications*. Jun 2018;10:121-130. doi:10.1016/j.conctc.2018.02.002
31. Dayan D, Kuriansky J, Abu-Abeid S. Weight Regain Following Roux-en-Y Gastric Bypass: Etiology and Surgical Treatment. *The Israel Medical Association journal : IMAJ*. Dec 2019;12(21):823-828.

ÍNDICE REMISSIVO

A

Acreditação hospitalar 116, 117, 118, 126

Ansiedad 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250

Antidepressivos 15, 17, 47, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170

Assédio moral 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212

Automedicação 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24

Avaliação psicológica 33, 150, 156

B

Bariatric surgery 252, 261, 262

C

Cabelo 55, 185, 186, 187, 188, 190, 193, 194, 195, 197

Câncer 29, 30, 53, 54, 55, 56, 57, 58, 72, 73, 74, 75, 76, 77, 78, 79, 99, 100, 101, 102, 138, 139, 148, 221, 222, 223, 225, 227, 231, 232, 234, 235, 236, 237

Cirurgias estéticas 150, 153, 156

Coagulopatias 213, 215, 216, 217, 218, 219

Corpo líquido 150

Covid-19 6, 7, 70, 103, 104, 114, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126

Criança 80, 82, 87, 88, 90, 92, 93, 94, 108, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 155

D

Diabetes 1, 2, 3, 4, 5, 6, 7, 8, 41, 42, 44, 46, 48, 49, 50, 51, 252, 253, 259, 262, 271

Doença de Von Willebrand 213, 215

Doma clássica 238, 240, 241, 242, 247, 250

E

Educação física 103, 105, 106, 107, 111, 236

Ensino híbrido 103, 105, 114, 115

Epstein-Barr Vírus (EBV) 5, 98

Espiritualidade 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 212

Exercício físico 55, 221, 222, 225

F

Fator VIII 213, 214, 215, 217

Feridas 45, 46, 48, 49, 171, 172, 174

Fonoaudiologia 80, 81, 82, 87, 94, 95, 96

G

Gagueira 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96

Gestão Hospitalar 117

H

Histocompatibilidade 59, 61, 63

Humanização 26, 32, 35, 128, 129, 132, 135

I

Íliaco 154, 263

Infecções virais 98

M

Mama 53, 54, 55, 56, 57, 58, 72, 73, 74, 75, 76, 77, 78, 79, 138, 139, 140, 142, 143, 144, 145, 148, 224, 230, 231, 234

N

Neoplasia 53, 72, 73, 74, 138, 139, 140, 148, 222, 266, 267

Nutritional and metabolic diseases 252

O

Óleo de coco 185, 187, 188, 190, 193, 194, 195, 196, 197, 198, 199

Oncologia 34, 78, 222

P

Papilomavírus Humano (HPV) 98, 99

Paracoccidiodomicose 137, 138, 139, 140, 148, 149

P. brasiliensis 138, 139

Pé diabético 6, 41, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52

Pediatria 113, 128, 135

Pele 15, 44, 47, 48, 63, 74, 138, 144, 154, 156, 173, 174, 185, 186, 187, 188, 190, 192, 194, 195, 196, 197, 198, 199, 214, 220, 223, 224

Polineuropatia diabética 41

Polissacarídeo celulósico 172, 182

Puerpério 35, 37, 40

Q

Qualidade de vida 2, 25, 31, 32, 34, 41, 43, 44, 50, 53, 54, 55, 57, 59, 60, 68, 75, 80, 82, 93, 94, 112, 132, 160, 214, 215, 221, 225, 231, 232, 233, 235

S

Sarcoma 263, 265, 266, 267, 268, 269, 270

Saúde 1, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 40, 41, 42, 43, 44, 45, 46, 47, 49, 50, 51, 53, 54, 55, 56, 57, 58, 59, 60, 68, 69, 72, 73, 74, 75, 76, 77, 78, 79, 81, 95, 96, 98, 100, 102, 104, 106, 107, 109, 111, 112, 114, 116, 117, 118, 119, 122, 123, 124, 125, 126, 127, 128, 129, 130, 132, 133, 134, 135, 136, 150, 152, 153, 156, 157, 158, 159, 160, 161, 164, 168, 169, 170, 186, 198, 200, 202, 203, 204, 206, 208, 210, 211, 212, 213, 214, 215, 216, 218, 219, 221, 222, 223, 224, 226, 227, 234, 235, 236, 261, 271

Saúde do trabalhador 10, 202, 208, 210, 211

Saúde mental 14, 17, 23, 28, 31, 33, 35, 37, 40, 53, 55, 152, 158, 161, 208, 212

Saúde pública 12, 13, 22, 23, 34, 35, 36, 40, 41, 42, 57, 74, 77, 96, 119, 200, 222, 227

Severe obesity 251, 252, 253, 258, 259

Sistema Único de Saúde - SUS 56, 72, 73, 77, 78, 79, 125, 160, 234

T

Tipagem HLA 59, 61, 62, 64, 65, 67, 68

Trabalho 10, 11, 13, 15, 16, 17, 19, 22, 24, 25, 29, 33, 37, 41, 44, 48, 50, 56, 59, 60, 61, 64, 73, 80, 82, 83, 90, 94, 98, 107, 108, 114, 123, 127, 129, 133, 134, 152, 153, 154, 155, 171, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 219, 221, 227, 231, 232, 233

Transplante 59, 60, 61, 62, 63, 65, 66, 67, 68, 69, 225

Transtornos psicóticos 33, 35, 37

U

Úlcera diabética 41, 44

V

Violência 87, 89, 90, 91, 92, 94, 95, 96, 156, 200, 201, 202, 203, 204, 205, 206, 207, 210, 211

Y

Youtube 5, 7, 8

CIÊNCIAS DA SAÚDE:

Oferta, acesso e utilização 2



-  www.atenaeditora.com.br
-  contato@atenaeditora.com.br
-  @atenaeditora
-  www.facebook.com/atenaeditora.com.br

CIÊNCIAS DA SAÚDE:

Oferta, acesso e utilização 2



🌐 www.atenaeditora.com.br
✉ contato@atenaeditora.com.br
📷 @atenaeditora
📘 www.facebook.com/atenaeditora.com.br

Atena
Editora
Ano 2022