# Journal of Engineering Research

# ORGANIC MICROPOLLUTANTS IN THE DIFFERENT ENVIRONMENTAL COMPARTMENTS

Amado Enrique Navarro Frómeta

"Universidad Tecnológica de Izúcar de Matamoros". Izúcar de Matamoros, Puebla, Mexico https://orcid.org 0000-0002-4827-0670



All content in this magazine is licensed under a Creative Commons Attribution License. Attribution-Non-Commercial-Non-Derivatives 4.0 International (CC BY-NC-ND 4.0). Abstract: Organic micropollutants (OMPs), also called pollutants of emerging concern in certain circumstances, encompass a large group of chemical substances of natural or anthropogenic origin that, due to their molecular characteristics, present risks to human health. These include, among other chemicals, personal care products, additives, flame retardants, and food pharmaceuticals. In the light of the literature on the subject, the presence of OMPs in the different environmental compartments is addressed: air, water and soil. Results of the determinations, by gas chromatography coupled to mass spectrometry, of the concentrations of a large group of OMPs in the atmospheric particulate matter of the city of Puebla, river water, irrigation water and groundwater from the Nexapa river basin, Puebla and in Colombian soils are presented. environmental In these matrices the concentrations were in the ppb to ppt range. Keywords: Micropollutants, air, water, soil.

## INTRODUCTION

The full and effective realization of the right to a healthy environment is threatened by contamination due to the presence of chemical substances, many of which have harmful effects on biota and affect the normal performance of ecosystems (Metcalfe et al., 2022). Already in 2019, the United Nations Environment Program indicated that the chemical industry had prospects of doubling by 2030, observing a rapid increase in production and consumption in emerging economies (UNEP 2019). The foregoing, considering only the contamination of water and the work environment, caused the loss of around 2.2 million lives in 2019, a possibly underestimated figure (Fuller et al., 2022). Globally there are more than 350,000 chemical substances and their mixtures registered for their production and use. It is evident that we have not yet advanced sufficiently in the evaluation of its risks to human health, which can even include damage to children from prenatal exposures, requiring alternative in vivo, in vitro and in silico methods to determine its effects in shorter times (Brooks, 2020).

One of the anthropic threats (not well known by the general public) for ecosystems comes from the emission to the different environmental compartments of the wide range of chemical substances that, due to technological development, the depletion or insufficiency of certain natural products and population growth, among other factors, are used by man in his daily activities (Krishnakumar et al., 2022). Its denomination is diverse: if its action on organisms is considered, endocrine disrupting compounds and carcinogenic substances are considered; based on their recent detection, attention and/or lack of regulation they are known as emerging pollutants or pollutants of emerging concern (EPs); Since they are found at a trace level in the environment, they are also called organic micropollutants, OMPs, a denomination to which this work adheres. The metabolites resulting from their transformations must also be considered, sometimes more dangerous than the starting compounds (Daughton, 2005; Libralato et al., 2020; Reyes et al., 2021). Being anthropogenic contaminants, they are found in all spheres of our planet and even in food, accumulating and biomagnifying in food chains (Krishnakumar et al., 2022; Das et al., 2023; Li et al., 2023). Events that affect the human species such as climate change and the recent pandemic affect the increase in its presence, for example, antidepressant drugs and other EPs (Castillo-Zacarías et al., 2021; Picó & Barceló, 2023). The most worrying thing is that we have not learned much about the impact of contamination by chemical substances on the

environment and on living beings, including the human species and, although there is some progress in knowledge and legislation on the most dangerous substances (called priority pollutants), legislation in Latin America on the regulation of EPs and OMPs is practically non-existent (Jacobo-Marín & Santacruz de León, 2021).

The objective of this work is to assess the information from the periodic literature on the presence of OMPs in the atmosphere, water and soil and to present the results of some experimental determinations of the same. This is intended to contribute to consolidating the social perception of these substances present in the environment.

## **OMPs AS CONTAMINANTS**

Table 1 shows a general classification of the most relevant OMPs (Navarro, 2019). Its structure in many cases favors its hydrophilicity and high solubility in water, which is why they can reach the cellular level when penetrating living organisms. As can be seen, its toxicology is very varied, including estrogenic and carcinogenic substances, among others, even reporting problems in fetal development due to maternal exposure to them (Hiranmai & Kamaraj, 2021; Sahoo et al., 2021; Cao et al., 2022).

Although they are not properly OMPs, it is essential to mention microplastics, which are unquestionably EPs. In addition to their wide presence on the planet, as well as the dangers they entail for biota, including their transfer to the food chain with humans as the upper link, they can contribute to the transport of OMPs that can be absorbed into them (Bhatt et al., 2021; Gola et al., 2021; Jain et al., 2021; Martin et al., 2023; Picó & Barceló, 2023).

For the determination of OMPs, various analytical techniques and bioassays are used, an aspect that is not discussed in this work, being the chromatographic methods coupled with mass spectrometry the most widespread. Most of the methods concern the determination of OMPs in aqueous matrices. As in any other chemical analysis, the correct preparation of the sample is decisive (di Paolo et al., 2016; Pena-Pereira et al., 2021; Hajeb et al., 2022; Chen L. et al., 2023).

Regarding the technologies for their removal, they have been more widely discussed and described for the aqueous medium, since it is the fundamental means of disposal of OMPs. It has also been reported that traditional wastewater treatment methods are not very effective for its removal, with advanced treatments, still relatively expensive, and nature-based systems such as phytoremediation techniques and treatment wetlands being more effective (Zhang, 2019; Lee et al., 2021; Shahid et al., 2021; Ilyas, 2021; Shi et al., 2023).

# **OMPS IN THE ATMOSPHERE**

OMPs and other EPs, such as antibioticresistant bacteria and genes, enter the atmosphere through volatilization, aerosol and bioaerosol formation, or diffusive exchange processes from their emission from various sources. Once in the atmosphere, they can be transported even long distances from emission sources. Its presence is more worrying in urban environments and due to the exchange between interiors and exteriors and its concentration in rooms and workplaces can be higher than in the exterior, for which reason inhalation is the most important route of exposure (Chen et al., 2022; Gwenzi, 2022; Simões et al., 2022). For its study, active or passive sampling techniques are used for both air and atmospheric particulate matter (PM). This way, OMPs belonging to practically all the types shown in Table 1 and volatile organic compounds (VOCs), of different types, have been detected and quantified in concentrations ranging from tenths of pg

Family	Example	Use/application	Toxicological effects	
Drugs	ethinyl estradiol	oral contraceptive	fish feminization	
personal care products (PCPs)	galaxolide	colonies and detergents	inhibitor of multixenobiotic resistance transporters	
perfluorinated compounds	perfluorooctanoic acid	non-stick pans with Teflon	cellular toxicity and possible human carcinogen	
surfactants	octylphenol	industrial detergents	fish feminization	
polymer additives	bisphenol A	plastic production	fish feminization	
polar pesticides	terbutylazine	herbicide	classified as possible carcinogen (group 2B)	
drugs of abuse	amphetamine	medical applications	hepatotoxic in fish	
flame retardants	polybrominated diphenyl ether	plastics and foams	impact on the development, growth and reproduction of crustaceans	
nanoparticles	Fullerenes	nanomaterials	possible cell damage due to oxidation	

Table 1. Classification of the most widely distributed OMPs (Navarro, 2019).

	Average	Median	Minimum	Maximum	Standard deviation	Standard error of the mean
DCB	0.179	0.182	0.024	0.708	0.152	0.026
ТСВ	0.371	0.313	0.020	1.405	0.329	0.056
DM3Oc	0.081	0.033	0.001	0.666	0.133	0.024
DMAcPhen	0.143	0.101	0.014	0.632	0.138	0.024
BzPhen	0.831	0.725	0.338	1.906	0.355	0.061
MDHJ	0.681	0.573	0.301	2.094	0.426	0.073
Gal	0.256	0.235	0.124	0.704	0.110	0.019
Ton	0.140	0.127	0.079	0.296	0.057	0.010
OctBenz	0.219	0.200	0.031	0.509	0.108	0.019
DOctSeb	0.278	0.138	0.019	1.062	0.296	0.050

Table 2. Some OMPs present in the atmosphere of the city of Puebla (ng m-3).

Note: DCB – dichlorobenzenes, TCB – 1,2,4-trichlorobenzene, DM3Oc – 2,3-dimethyl-3-octanol, DMAcPhen – 2',4'-acetophenone, BzPhen – benzophenone, MDHJ – Methyldihydrojasmonate; Gal-Galaxolide; Ton – Tonalide; OctBenz – octylbenzoate, DoctSeb – dioctyl sebasate. m-3 to hundreds of ng m-3 (Barroso et al., 2019; Hsiao et al., 2019; Enyoh et al., 2020; Robichaud 2020; Kim et al., 202 1).

During the years 2009 and 2010, samples of PM smaller than 10 µm were analyzed, coming from 3 stations of the Environmental Monitoring Network of the City of Puebla. Table 2 shows the descriptive statistics of some of the OMPs identified and determined. As can be seen, VOCs, plasticizers and, as expected, the musk-type fragrances with a broad spectrum of use MDHJ, Gal and Ton were detected and quantified. Some alkylamides possibly from biomass burning were detected but not quantified. Temporal patterns of some of the substances studied did not show consistency with seasonal temperature, suggesting that other factors influence pollutant concentrations beyond volatilization (Navarro, 2013).

# OMPs IN THE AQUATIC ENVIRONMENT

Since freshwater bodies are the destination of many treated and untreated wastewater discharges, most of the studies on the presence of OMPs have been aimed at determining their concentrations in surface streams, lakes and reservoirs, estuaries and, of course, drinking water supply sources. Given the possible migration routes due to point or diffuse contamination and their presence in the biota through trophic chains, their fate and bioaccumulation have also been of interest in many studies, a relevant aspect if we consider the consumption of aquatic species by humans (Alagan et al., 2023; Chen X. et al., 2023; Guo et al., 2023). It must be considered that water contamination by OMPs is not a static phenomenon and the lists and priorities regarding the OMPs to study are affected by some important variables: climate change; demographic changes, substitution of some chemicals with less problematic ones, new evidence on the toxicology and effects of OMPs, and technological development with the emergence of new applications of chemicals (Bunke et al., 2019, Yankui et al., 2019; Deere et al., 2021; Hatje et al., 2022).

In studies carried out in the upper part of the course of the Nexapa River in Puebla, Mexico, 398 OMPs were identified in the samples obtained in campaigns carried out in the years 1999 to 2013 (Figure 1). Of these, most are used in daily life as personal care products, cleaning products, food components, pharmaceuticals and pesticides. Of these, a group of 16 individual compounds and 3 families of compounds were quantified, observing the increase in concentration in dry season, at the point of transfer of 4 m3 per second of heavily contaminated water from the Atoyac River and after the most important populations as shown in figure 1B (Navarro et al., 2014). As can be seen in figure 1A, as a result of a principal component analysis carried out with the OMPs concentrations, four groups of samples are distinguished, made up of the SD and LF samples in dry conditions, those from those points in rain, and the samples from the rest of the points differentiated into dry and rain. The LM samples are closest to the vertex of the minimum values of the three axes. This way, the PCA summarizes the main spatiotemporal trends of OMPs concentrations observed in this study.

OMPs have been used as chemical markers to study the impact of wastewater on freshwater (Lim et al., 2017). Similarly, they can be used to assess the anthropogenic impact on coastal marine ecosystems, as shown in Figure 2, for the determination of alkylphenOMPs (AF) and coprostanol (COP) in waters and sediments of Cispatá Bay. As can be seen, the concentrations of COP in the sediments reflect the dragging of these by the Sinú River, while the AF reflect the incipient impact of human settlements located on the coastline (Navarro et al., 2016).

In the irrigation waters in the Atlixco-Izúcar valley, Puebla, part of which is derived from the Nexapa river, numerous OMPs were also detected (Navarro et al., 2017). Figure 3 shows the concentrations of these, compared with those present in irrigation water with similar characteristics from a region of Spain (Calderón-Preciado et al; 2011a). As it can be seen, the concentrations detected in Mexico are much higher than those determined in Spain. Unquestionably, this represents a greater risk of these contaminants passing through the food chain and reaching humans.

The impact of OMPs also reaches groundwater (Pradhan et al., 2023). In wells for irrigation in the Izúcar region, some OMPs were detected (CAF, GAL, TON, Sunscreen, MDHJ, AF and monoethoxylated alkylphenOMPs) in concentrations up to 13  $\mu$ g L<sup>-1</sup> (Navarro et al., 2019).

# **OMPs IN SOILS**

The main source of OMPs in agricultural soils is wastewater irrigation (Lyu et al., 2022; Minhas et al., 2022). Once present, both aerobic and anaerobic degradation processes occur, with greater speed than the former. Regardless of the degradation by microorganisms and other components of the soil biota, OMPs are absorbed by different routes by crops. Different factors influence this, such as plant species, the type of contaminants, microorganisms present in the soil, and environmental factors such as temperature and soil erosion. This shows the importance of controlling the levels of these contaminants in the soil to assess the risks due to it (Pullagurala et al., 2018; Biel-Maeso et al., 2019; García Valverde et al., 2021; Murrel et al., 2021).

The determination of some OMPs in Colombian agricultural soils irrigated with

water from the Sinú river, shows that the sum of the concentrations of 5 OMPs turned out to be 27.7 g ha-1 (Navarro et al., 2017), much higher than the sum of these same compounds in a soil from a region of Spain (Calderón Preciado et al., 2011b).

# CONCLUSIONS

are ubiquitous **OMPs** pollutants in environmental compartments, different generally having, as the main source, the disposal of personal care products, drugs, surfactants and other substances in domestic and municipal wastewater. In Latin America, more studies are needed to determine their levels in the different environmental matrices and legislation to begin regulating their levels in them and mitigate the risks due to the exposure of the population to these pollutants.

## REFERENCES

Alagan, M., Chandra Kishore, S., Perumal, S., Manoj, D., Raji, A., Kumar, R. S., Almansour, A. I., & Lee, Y. R. (2023). Narrative of hazardous chemicals in water: Its potential removal approach and health effects. *Chemosphere*, 335(139178), 139178. https://doi.org/10.1016/j.chemosphere.2023.139178

Barroso, P. J., Santos, J. L., Martín, J., Aparicio, I., & Alonso, E. (2019). *Emerging contaminants in the atmosphere: Analysis, occurrence and future challenges. Critical Reviews in Environmental Science and Technology,* 1–68. doi:10.1080/10643389.2018. 1540761

Bhatt, P., Pathak, V. M., Bagheri, A. R., & Bilal, M. (2021). *Microplastic contaminants in the aqueous environment, fate, toxicity consequences, and remediation strategies. Environmental Research, 200, 111762.* doi:10.1016/j.envres.2021.111762

Biel-Maeso, M., González-González, C., Lara-Martín, P. A., & Corada-Fernández, C. (2019). Sorption and degradation of contaminants of emerging concern in soils under aerobic and anaerobic conditions. Science of The Total Environment, 666, 662–671. doi:10.1016/j.scitotenv.2019.02.279

Brooks, B. W., Sabo-Attwood, T., Choi, K., Kim, S., Kostal, J., LaLone, C. A., ... Zhang, X. (2020). *Toxicology Advances for 21st Century Chemical Pollution. One Earth*, 2(4), 312–316. doi:10.1016/j.oneear.2020.04.007

Bunke, D., Moritz, S., Brack, W., Herráez, D. L., Posthuma, L., & Nuss, M. (2019). Developments in society and implications for emerging pollutants in the aquatic environment. Environmental Sciences Europe, 31(1). doi:10.1186/s12302-019-0213-1

Calderón-Preciado, D., Matamoros, V., & Bayona, J. M. (2011a). Occurrence and potential crop uptake of emerging contaminants and related compounds in an agricultural irrigation network. *The Science of the Total Environment*, 412–413, 14–19. doi: 10.1016/j.scitotenv.2011.09.057

Calderón-Preciado, D., Jiménez-Cartagena, C., Matamoros, V., & Bayona, J. M. (2011b). Screening of 47 organic microcontaminants in agricultural irrigation waters and their soil loading. *Water Research*, 45(1), 221–231.

Cao, Y., Chen, Z., Zhang, M., Shi, L., Qin, S., Lv, D., Li, D., Ma, L. and Zhang, Y. (2021). Maternal exposure to bisphenol A induces fetal growth restriction via upregulating the expression of estrogen receptors. *Chemosphere*, 287, p.132244. doi.org/10.1016/j. chemosphere.2021.132244

Castillo-Zacarías, C., Barocio, M. E., Hidalgo-Vázquez, E., Sosa-Hernández, J. E., Parra-Arroyo, L., López-Pacheco, I. Y., Barceló, D., Iqbal, H. N. M., & Parra-Saldívar, R. (2021). Antidepressant drugs as emerging contaminants: Occurrence in urban and non-urban waters and analytical methods for their detection. *The Science of the Total Environment*, 757(143722), 143722. https://doi. org/10.1016/j.scitotenv.2020.143722

Chen, L., Yan, X., Zhou, X., Peng, P., Sun, Q., & Zhao, F. (2023). Advances in the on-line solid-phase extraction-liquid chromatography-mass spectrometry analysis of emerging organic contaminants. *Trends in Analytical Chemistry: TRAC*, *160*(116976), 116976. https://doi.org/10.1016/j.trac.2023.116976

Chen, P., Guo, X., & Li, F. (2022). Antibiotic resistance genes in bioaerosOMPs: Emerging, non-ignorable and pernicious pollutants. *Journal of Cleaner Production*, 348(131094), 131094. https://doi.org/10.1016/j.jclepro.2022.131094

Chen, X., Qadeer, A., Liu, M., Deng, L., Zhou, P., Mwizerwa, I. T., Liu, S., Ajmal, Z., Xingru, Z., & Jiang, X. (2023). Bioaccumulation of emerging contaminants in aquatic biota: PFAS as a case study. In *Emerging Aquatic Contaminants* (pp. 347–374). Elsevier.

Das, C., Lincy G., M. V., Shafi, T., Dubey, B. K., Ghangrekar, M. M., & Chowdhury, S. (2023). Ecological effects of emerging contaminants: Ecotoxicity and relative environmental risk. In *Emerging Aquatic Contaminants* (pp. 375–404). Elsevier.

Daughton CG. (2005). "Emerging" chemicals as pollutants in the environment: a 21st century perspective". *Renewable Resources Journal* 23.4: 6.

Deere, J. R., Streets, S., Jankowski, M. D., Ferrey, M., Chenaux-Ibrahim, Y., Convertino, M., ... Wolf, T. M. (2021). A chemical prioritization process: Applications to contaminants of emerging concern in freshwater ecosystems (Phase I). Science of The Total Environment, 772, 146030. doi:10.1016/j.scitotenv.2021.146030

Di Paolo, C., Ottermanns, R., Keiter, S., Ait-Aissa, S., Bluhm, K., Brack, W., ... Hollert, H. (2016). *Bioassay battery interlaboratory investigation of emerging contaminants in spiked water extracts – Towards the implementation of bioanalytical monitoring toOMPs in water quality assessment and monitoring. Water Research*, 104, 473–484. doi:10.1016/j.watres.2016.08.018

Enyoh, C. E., Verla, A. W., Qingyue, W., Ohiagu, F. O., Chowdhury, A. H., Enyoh, E. C., ... Chinwendu, U. P. (2020). An overview of emerging pollutants in air: Method of analysis and potential public health concern from human environmental exposure. Trends in Environmental Analytical Chemistry, e00107. https://doi.org/10.1016/j.teac.2020.e00107

Fuller, R., Landrigan, P. J., Balakrishnan, K., Bathan, G., Bose-O'Reilly, S., Brauer, M., Caravanos, J., Chiles, T., Cohen, A., Corra, L., Cropper, M., Ferraro, G., Hanna, J., Hanrahan, D., Hu, H., Hunter, D., Janata, G., Kupka, R., Lanphear, B., ... Yan, C. (2022).
Pollution and health: A progress update. *The Lancet Planetary Health*, 6(6). https://doi.org/10.1016/s2542-5196(22)00090-0

Gola, D., Kumar Tyagi, P., Arya, A., Chauhan, N., Agarwal, M., Singh, S. K., & Gola, S. (2021). The impact of microplastics on marine environment: A review. Environmental Nanotechnology, Monitoring & Management, 16, 100552. doi:10.1016/j. enmm.2021.100552

Guo, J., Tu, K., Chou, L., Zhang, Y., Wei, S., Zhang, X., Yu, H., & Shi, W. (2023). Deep mining of reported emerging contaminants in China's surface water in the past decade: Exposure, ecological effects and risk assessment. *Water Research*, 243(120318), 120318. https://doi.org/10.1016/j.watres.2023.120318

Gwenzi, W. (2022). Air-borne emerging contaminants: An under-studied reservoir and a potential health risk? In *Emerging Contaminants in the Terrestrial-Aquatic-Atmosphere Continuum*: (pp. 139–150). Elsevier.

Hajeb, P., Zhu, L., Bossi, R. and Vorkamp, K. (2022). Sample preparation techniques for suspect and non-target screening of emerging contaminants. *Chemosphere*, 287, part 3, p.132306. https://doi.org/10.1016/j.chemosphere.2021.132306

Hatje, V., Sarin, M., Sander, S. G., Omanović, D., Ramachandran, P., Völker, C., Barra, R. O., & Tagliabue, A. (2022). Emergent interactive effects of climate change and contaminants in coastal and ocean ecosystems. *Frontiers in Marine Science*, 9. https://doi.org/10.3389/fmars.2022.936109

Hiranmai, R. & Kamaraj, M. (2021). Occurrence, fate, and toxicity of emerging contaminants in a diverse ecosystem. Physical Sciences Reviews, (), 000010151520210054. https://doi.org/10.1515/psr-2021-0054

Hsiao, T.-C., Lin, A. Y.-C., Lien, W.-C., & Lin, Y.-C. (2019). Size Distribution, Biological Characteristics and Emerging Contaminants of AerosOMPs Emitted from an Urban Wastewater Treatment Plant. Journal of Hazardous Materials, 121809. doi:10.1016/j.jhazmat.2019.121809

Ilyas, H. (2021). Emerging Organic Contaminants Removal by Constructed Wetlands: Comprehensive Meta-analysis and Development of Decision Support ToOMPs. Thèse de doctorat de Sciences de la Terre et de l'Environnement. Université de Paris.

Jacobo-Marín, D., & Santacruz de León, G. (2021). Contaminantes emergentes en el agua: Regulación en México, principio precautorio y perspectiva comparada. *Revista De Derecho Ambiental*, 1(15), pp. 51–75. https://doi.org/10.5354/0719-4633.2021.57414

Jain, S., Mishra, D., & Khare, P. (2021). Microplastics as an Emerging Contaminant in Environment: Occurrence, Distribution, and Management Strategy. Management of Contaminants of Emerging Concern (CEC) in Environment, 281–299. doi:10.1016/ b978-0-12-822263-8.00010-5

Kim, C., Choi, D., Lee, Y. G., & Kim, K. (2021). *Diagnosis of indoor air contaminants in a daycare center using a long-term monitoring. Building and Environment, 204, 108124.* doi:10.1016/j.buildenv.2021.10812

Krishnakumar, S., Singh, D. S., Godson, P. S., & Thanga, S. G. (2022). Emerging pollutants: Impact on environment, management, and challenges. *Environmental Science and Pollution Research*, *29*(48), 72309–72311. https://doi.org/10.1007/s11356-022-22859-3

Lee, B. C. Y., Lim, F. Y., Loh, W. H., Ong, S. L., & Hu, J. (2021). Emerging Contaminants: An Overview of Recent Trends for Their Treatment and Management Using Light-Driven Processes. *Water*, *13*(17), 2340. doi:10.3390/w13172340

Li, X., Liu, Y., Yin, Y., Wang, P., & Su, X. (2023). Occurrence of some legacy and emerging contaminants in feed and food and their ranking priorities for human exposure. *Chemosphere*, *321*(138117), 138117. https://doi.org/10.1016/j.chemosphere.2023.138117

Libralato, G., Freitas, R., Buttino, I., Arukwe, A., & Della Torre, C. (2020). Special issue on challenges in emerging environmental contaminants CEEC19. Environmental Science and Pollution Research. doi:10.1007/s11356-020-09539-w

Lim, F., Ong, S., & Hu, J. (2017). Recent Advances in the Use of Chemical Markers for Tracing Wastewater Contamination in Aquatic Environment: A Review. Water, 9(2), 143. doi:10.3390/w9020143

Lyu, S., Wu, L., Wen, X., Wang, J., & Chen, W. (2022). Effects of reclaimed wastewater irrigation on soil-crop systems in China: A review. *The Science of the total environment*, *813*, 152531. https://doi.org/10.1016/j.scitotenv.2021.152531

Martín, J., Santos, J. L., Aparicio, I., & Alonso, E. (2022). Microplastics and associated emerging contaminants in the environment: Analysis, sorption mechanisms and effects of co-exposure. *Trends in Environmental Analytical Chemistry*, 35(e00170), e00170. https://doi.org/10.1016/j.teac.2022.e00170

Metcalfe, C. D., Bayen, S., Desrosiers, M., Muñoz, G., Sauvé, S., & Yargeau, V. (2022). An introduction to the sources, fate, occurrence and effects of endocrine disrupting chemicals released into the environment. *Environmental Research*, 207(112658), 112658. https://doi.org/10.1016/j.envres.2021.112658

Minhas, P. S., Saha, J. K., Dotaniya, M. L., Sarkar, A., & Saha, M. (2022). Wastewater irrigation in India: Current status, impacts and response options. *The Science of the total environment*, 808, 152001. https://doi.org/10.1016/j.scitotenv.2021.152001

Murrell, K. A., Teehan, P. D., & Dorman, F. L. (2021). *Determination of contaminants of emerging concern and their transformation products in treated-wastewater irrigated soil and corn. Chemosphere*, 281, 130735. doi:10.1016/j.chemosphere.2021.130735

Navarro A. (2013). Anthropic Signatures in the Urban Particulate of the City of Puebla, México, Proceedings of a meeting held 17-21 March 2013, Philadelphia, Pennsylvania, USA. Abstracts. PITTCON Conference and Expo 2013. Curran Associates, Inc. (Oct 2014).

Navarro A. (2019). Consideraciones sobre el estudio de los microcontaminantes orgánicos en México En Memorias del Noveno Minisimposio Internacional Sobre Remoción De Contaminantes De Aguas, Atmósfera Y Suelos. Eds. Garrido-Mora A., Durán-Domínguez-de-Bazúa M.C. Pp. 143-152. Villahermosa, Tabasco, México. Disponible en https://ambiental.unam.mx/ albunimagenes/Libro-e-Book%202019%20IX%20MSI-9th%20IMS.pdf

Navarro A., Herrera J., Marrugo J., Bayona J., Morales L. (2014). Microcontaminantes orgánicos en los ríos de México: El caso del río Nexapa. En M. Ramos., V. Aguilera., (eds.) .Ciencias de la Ingeniería y Tecnología, Handbook T-IV. -©ECORFAN- Valle de Santiago, Guanajuato, pp. 49-63.

Navarro A., Marrugo J.L., Bayona J. (2017). Selected organic micropollutants in agricultural environments of México and Colombia: a comparison with reported levels in Spain. Book of proceedings II-International Caparica Conference on Pollutant Toxic Ions and Molecules, p. 129. Lisboa, Portugal 6 al 9 de noviembre de 2017. ISBN 978-989-99639-8-6.

Navarro A.E., Marrugo J.L., Burgos S.M., Morales L., Consuegra A., Herrera J.A. (2016). Huellas antrópicas en la Bahía de Cispatá, Colombia: un estudio de línea base de microcontaminantes orgánicos. Memorias del VI Congreso Iberoamericano de Química Analítica, pp. 84-89. Red Iberoamericana de Química Analítica, Universidad Autónoma del Estado de México. Cancún, Q. Roo.

Navarro-Frómeta A.E., Herrera-López H., Castro-Bravo C. (2019). Irrigation Water Challenges: A Study Case in the State of Puebla, Mexico. In: Water Availability and Management in Mexico. Editors: Elena María Otazo-Sánchez, Amado Enrique Navarro-Frómeta, Vijay P. Singh. Springer Nature, Switzerland, AG 2020. Pages 249-265.

Pena-Pereira, F., Bendicho, C., Pavlović, D. M., Martín-Esteban, A., Díaz-Álvarez, M., Pan, Y., ... Psillakis, E. (2020). *Miniaturized analytical methods for determination of environmental contaminants of emerging concern – A Review. Analytica Chimica Acta.* doi:10.1016/j.aca.2020.11.040

Picó, Y., & Barceló, D. (2023). Microplastics and other emerging contaminants in the environment after COVID-19 pandemic: The need of global reconnaissance studies. *Current Opinion in Environmental Science & Health*, 33(100468), 100468. https://doi. org/10.1016/j.coesh.2023.100468

Pradhan, B., Chand, S., Chand, S., Rout, P. R., & Naik, S. K. (2023). Emerging groundwater contaminants: A comprehensive review on their health hazards and Remediation Technologies. *Groundwater for Sustainable Development*, *20*, 100868. https://doi.org/10.1016/j.gsd.2022.100868

Pullagurala, V. L. R., Rawat, S., Adisa, I. O., Hernandez-Viezcas, J. A., Peralta-Videa, J. R., & Gardea-Torresdey, J. L. (2018). *Plant uptake and translocation of contaminants of emerging concern in soil. Science of The Total Environment, 636, 1585–1596.* doi:10.1016/j.scitotenv.2018.04.375

Reyes, N.J.D.G.; Geronimo, F.K.F.; Yano, K.A.V.; Guerra, H.B.; Kim, L.-H. (2021). Pharmaceutical and Personal Care Products in Different Matrices: Occurrence, Pathways, and Treatment Processes. Water, 13, 1159. https://doi.org/10.3390/ w13091159

Robichaud, A. (2020). An overview of selected emerging outdoor airborne pollutants and air quality issues: the need to reduce uncertainty about environmental and human impacts. Journal of the Air & Waste Management Association. doi:10.1080/109622 47.2020.1723738

Sahoo, T., Sahu, J., Panda, J., Nayak, P., Soren, S., Sahoo, S. and Sahu, R. (2020). Carcinogenic Nature of Emerging Contaminants: Havoc for Present and Gateway of Unhealthy Future. *Contaminants in Drinking and Wastewater Sources*, pp.159-175. https://doi.org/10.1007/978-981-15-4599-3\_7

Shahid, M. K., Kashif, A., Fuwad, A., & Choi, Y. (2021). Current advances in treatment technologies for removal of emerging contaminants from water – A critical review. Coordination Chemistry Reviews, 442, 213993. doi:10.1016/j.ccr.2021.213993

Shi, Q., Kaur, P., & Gan, J. (2023). Harnessing the potential of phytoremediation for mitigating the risk of emerging contaminants. *Current Opinion in Environmental Science & Health*, 32(100448), 100448. https://doi.org/10.1016/j.coesh.2023.100448

Simões, M., Janssen, N., Heederik, D. J. J., Smit, L. A. M., Vermeulen, R., & Huss, A. (2022). Residential proximity to livestock animals and mortality from respiratory diseases in The Netherlands: A prospective census-based cohort study. *Environment International*, *161*(107140), 107140. https://doi.org/10.1016/j.envint.2022.107140

UNEP, 2019. United Nations Environment Program. Global Chemicals Outlook II: *From Legacies to Innovative Solutions: Implementing the 2030 Agenda for Sustainable Development*. ISBN: 978-92-807-3745-5. Disponible en: https://wedocs.unep.org/bitstream/handle/20.500.11822/28113/GCOII.pdf?sequence=1&isAllowed=y

Yankui, T., Yin, M., Yang, W., Li, H., Zhong, Y., Mo, L., ... Sun, X. (2019). *Emerging pollutants in water environment: occurrence, monitoring, fate, and risk assessment. Water Environment Research.* doi:10.1002/wer.1163

Zhang, D. (2019). The role of microorganisms in the removal of pharmaceutical and personal care products. Pharmaceuticals and Personal Care Products: Waste Management and Treatment Technology, 341–382. doi:10.1016/b978-0-12-816189-0.00015-9




Figure 1. Main trends of OMPs concentrations in the Nexapa River. A – PCA results; B – Concentration of triclosan at the sampling points. Location of sampling points: 1 – LM (Los Molinos); 2 – SD (Santo Domingo Atoyatempan), after the transfer of water from the Atoyac; 3 – LF (Las Fajanas, after the city of Atlixco); 4 – Al (Alchichica); 5 – Puc (Puctla before discharge from the treatment plant); 6 – Pu2 (Puctla after discharge from the plant).




Figure 2. COP and AF concentrations in samples from Cispatá Bay (source Navarro et al., 2016).




Figure 3. OMPs concentrations in irrigation waters from Mexico and Spain (adapted from Navarro-Frómeta et al., 2017); MDHJ – methyl-dihydrojasmonate, CAF – caffeine, BHT – butylated hydroxytoluene, TCS – triclosan, AP – alkylphenols- NAP - naproxen